



GOVERNMENT OF KERALA

# DISTRICT SURVEY REPORT OF MINOR MINERALS (EXCEPT RIVER SAND)

## KOLLAM DISTRICT

Prepared as per  
Environment Impact Assessment (EIA) Notification, 2006 issued  
under Environment (Protection) Act 1986  
by

DEPARTMENT OF MINING AND GEOLOGY  
[www.dmg.kerala.gov.in](http://www.dmg.kerala.gov.in)

November, 2016  
Thiruvananthapuram

# Table of Contents

	Page No.
1 Introduction .....	3
2 Drainage and Irrigation .....	5
3 Rainfall and climate .....	6
4 Meteorological Parameters .....	6
4.1 Temperature .....	6
4.2 Relative Humidity .....	6
4.3 Evaporation .....	6
4.4 Sunshine Hours.....	6
4.5 Wind.....	6
4.6 Potential Evapotranspiration (PET) .....	7
5 Geology .....	7
6 Geomorphology .....	10
7 Soil types .....	10
8 Land use .....	10
9 Forest .....	11
10 Groundwater scenario .....	11
11 Natural hazards.....	17
12 Mineral Resources .....	17
12.1 Major minerals .....	17
12.2 Minor Minerals.....	17
12.2.1 Ordinary Earth.....	17
12.2.2 Ordinary Clay (tile/brick clay) .....	22
12.2.3 Ordinary Sand .....	24

12.2.4	Laterite.....	25
12.2.5	Granite Dimension Stone and Granite (building stone) .....	27
13	Details of minor mineral concessions and revenue collection .....	28

### **List of Figures**

Figure 1: Geology and mineral resources of Kollam.

Figure 2: Geomorphology and geohydrology of Kollam.

Figure 3: Land use of Kollam.

Figure 4: Geotechnical characteristics and natural hazards map of Kollam.

Figure 5: Soil map of Kollam

### **List of Tables**

Table1: Details of revenue collection for the period 2013-'14, 2014-'15 and 2015-'16

Table 2a: List of Quarrying Lease granted for Granite building stone

Table 2b: List of Quarrying Permit granted under CRPS for Granite Building Stone (2015-'16)

Table 2c: List of Quarrying Permit granted under CRPS for Granite Building Stone (2016-'17)

Table 2d: List of short term permits granted for Granite Building Stone (2016-'17)

Table 2e: List of Quarrying Permit granted under CRPS for Laterite Building Stone during 2015-'16

Table 2f: List of Quarrying Permit granted under CRPS for Laterite Building Stone during 2016-'17

Table 2g: List of Registered Metal Crusher Units - RMCU (2016-'17)

Annexure 1. Geology of Kerala

# DISTRICT SURVEY REPORT OF MINOR MINERALS

## KOLLAM DISTRICT


**(This report is to be submitted along with application for Environmental Clearance (EC) for mining of all minor minerals except river sand)**

### 1 Introduction

Kollam or Quilon, an old sea port town on the Arabian coast, stands on the Ashtamudi lake. Kollam, the erstwhile Desinganadu, had a sustained commercial reputation from the days of the Phoenicians and the Romans. Fed by the Chinese trade, it was regarded by Ibn Batuta, as one of the five ports, which he had seen in the course of his travels during a period of twenty four years, in the 14th century. Kollam District which is a veritable Kerala in miniature is gifted with unique representative features - sea, lakes, plains, mountains, rivers, streams, backwaters, forest, vast green fields and tropical crop of every variety both food crop and cash crop, so called '*The Gods Own Capital*'. Kollam District is located on the southwest part of Kerala State and extends from Lakshadweep Sea to the Western Ghats and is bordered by Trivandrum district on the South and Alleppey and Pathanamthitta districts in the North and Tirunelveli district of Tamil Nadu State in the East and Lakshadweep Sea in the west. It has a maximum length of 75 kms in the E-W direction and maximum width of 45 kms in the N-S direction. It lies between North latitudes 8° 45' and 9° 07' and East longitudes 76° 29' and 77° 17'. It has a geographical area of 2412 sq. km. and falls in parts of Survey of India Toposheets 58C, D, G and H.

The Quilon district is divided both on geographical and functional basis for purposes of general administration. Geographically it is divided into Revenue Divisions. Kollam is administratively divided into 6 taluks. They are Kollam, Karunagappally, Kunnathur, Kottarakkara, Punalur and Pathanapuram, which are subdivided into 104 villages.

In 2011, Kollam had population of 2,635,375 of which male and female were 1,246,968 and 1,388,407 respectively. In 2001 census, Kollam had a population of 2,585,208 of which males were 1,249,621 and remaining 1,335,587 were females. The literacy rate of the district is 92.66.



INDIA  
KERALA  
**KOLLAM DISTRICT**



NUMBER OF TALUKS.....	: 6
STATUTORY TOWNS .....	: 3
CENSUS TOWNS .....	: 24
OUT GROWTHS .....	: 2
VILLAGES .....	: 69
TOTAL AREA OF THE DISTRICT .....	: 2491 Sq. Km.

District headquarters is also the Taluk headquarters

BOUNDARY, STATE	—————
" DISTRICT	—————
" TALUK	—————
HEADQUARTERS - DISTRICT, TALUK	●
NATIONAL HIGHWAY	—————
STATE HIGHWAY	—————
OTHER IMPORTANT ROADS	—————
RAILWAY LINE, BROAD GAUGE	—————
RIVER AND STREAM	~~~~~
STATUTORY TOWN / CENSUS TOWN & OUT GROWTH	○

## 2 Drainage and Irrigation

The district is drained by three west flowing rivers , viz *Achenkovil*, *Kallada* and *Ithikara*, originating in the eastern hilly region. These rivers together with their tributaries exhibit dendritic pattern of drainage.

The Ithikara basin has its elevation north of *Madathara* (271 m amsl) on the eastern side and slopes down to sea level west of Mayyanad. The Ithikara river originates from the *Madatharaikunnu* hills, south west of Kulathupuzha and drains into the Paravoor backwaters near Meenad. Ithikara river is a fourth order stream with a slope of 8.2 m/km. The length of the river is 56 km and the drainage area is 779 km<sup>2</sup>.

The Kallada river basin has its highest elevation at *Karimalaikodkal* (1763 m amsl) on the eastern side and reaches almost sea level west of Karunagapally. The river originating from the Western Ghats drains into Ashtamudi backwaters near Kollam. The length of the river is 121 km and drainage area is 1996 km<sup>2</sup>. Kallada river is a fifth order stream with a gradient of 12.6 m/km.

The Achenkovil river originates from the Western Ghats and covers a basin area of 1484 km<sup>2</sup> and the main channel length is 128 km. The River joins Pamba river at Veeyapuram and finally debouches into the Vembanad lake. The Achankovil river is set in a well known shear zone demarcating the boundary between Kerala Khondalite Belt and charnockites of Southern Granulite terrain.

The district is blessed with the largest fresh water lake in the State namely the Sasthamkotta lake and is one of the resources which caters to the drinking water needs of Kollam district. The lake occupies 440 hectares and the catchment area of the lake is 1269 sq km. Other major lakes (Kayals) in the district are - Ashtamudi Kayal (6424 ha) and Paravoor Kayal (662 ha).

The irrigation facilities in the district are limited. The major irrigation scheme is Kallada irrigation project and the target fixed for it was 61630 ha of land and 92806 ha of crops. There are also minor irrigation schemes through which 1500 ha of land is being irrigated. Among source of irrigation, ground water is the principal source of irrigation accounting for about 47% of the area under irrigation and the rest by lift and other methods of irrigation.

### **3 Rainfall and climate**

The district receives an annual average rainfall of about 2428 mm. The Southwest monsoon from June to September contributes nearly 55% of the total annual rainfall. The Northeast monsoon season from October to December contributes about 24% and the balance 21% is received during the month of January to May as pre-monsoon showers. Out of the total 119 rainy days, about 70 rainy days occur during the southwest monsoon season.

### **4 Meteorological Parameters**

#### **4.1 Temperature**

The temperature is more during the months of March to May and is less during December and January. The average mean monthly maximum temperature ranges from 29.9 to 36.4°C and minimum temperature ranges from 19.4 to 23.8°C.

#### **4.2 Relative Humidity**

The Relative humidity is higher during the monsoon period and all through the year it is higher during the morning hours.

#### **4.3 Evaporation**

Evaporation is more during summer months of January to April and it is low during the rainy months May to August. The maximum rate of 4.8 mm per day is recorded in March and the lowest rate of 2.6 mm is recorded during July.

#### **4.4 Sunshine Hours**

Sunshine ranges from 4.3 to 9.7 hours/day. Maximum sunshine is during the month of February. The months of June to August record the minimum sunshine due to the cloudy sky. Generally good sunshine hours are recorded in the months of November to May.

#### **4.5 Wind**

The wind speed ranges from 1.3 to 2.1 km/hour. The wind speed is high during the months of March to June and less during the months of September to December.



## 4.6 Potential Evapotranspiration (PET)

PET values are lower than the monthly rainfall during the month of May to October indicating water surplus for possible recharge into groundwater regime during these months. The monthly PET ranges from 119.3 to 177 mm.

## 5 Geology

The district can be broadly divided into three geological provinces – the westernmost Quaternary alluvial deposits followed by a narrow N-S zone of late Tertiary sediments and the easternmost Precambrian metamorphic.

The Precambrian metamorphic are represented by Khondalite, Charnockite and Migmatitic groups. They are intruded by younger basic dykes and overlain by Tertiary and Quaternary sediments.

High grade metamorphic rocks of Khondalite Group include calc-granulite, quartzite and garnet-biotite-sillimanite gneiss with or without graphite. Thin lenticular bands of calc-granulite occur within charnockite and migmatite. The Khondalite paragneiss tends to occur as linear bodies towards the middle and western part of the district.

The Charnockite Group consists of pyroxene granulite, cordierite gneiss and hypersthene-hornblende granite gneiss (charnockite). It mostly occurs as concordant bands and lenses of varied dimensions in Khondalite and migmatite with a diffused contact. It grades into gneiss. Generally, it is garnetiferous near the contact with the gneiss. The rock shows granoblastic texture and is mostly intermediate to acidic. Pyroxene granulite occurs as thin, discordant, lenticular patches, within migmatite, and is concordant with the para gneiss. Cordierite gneiss is found as impersistent bands and lenses within garnet-biotite gneiss and is confined to the contact with gneiss and charnockite. It displays xenoblastic gneissose texture and consists of varying proportions of cordierite, plagioclase, microperthite, quartz, biotite, hypersthene, garnet and hornblende. Near Punalur, there is a small body of dunite.

The Migmatite Complex comprising garnet-biotite gneiss and quartzo-feldspathic gneiss are the major rock units of the area and they are traversed by the pegmatite and aplite veins. The rocks of the Migmatite Complex are widely distributed and interlayered with rocks of Charnockite Group. Garnet-biotite gneiss has a larger areal distribution and is characterised by the presence of biotite foliae and concentration of



garnet in layers. Bands and lenses of quartzo-feldspathic gneiss occur within it. Granite gneiss of Peninsular Gneissic Complex occurs as small lenses towards the east. All the older rocks are intruded by basic intrusive of doleritic composition having a general NW-SE trend.

Towards west, the rocks of Archaean age are uncoformably overlain by sedimentary rocks of Mio-Pliocene age. Two distinct sequence of sediments are noticeable. A lower marine sequence (Quilon Formation) represented by fossiliferous limestone and marl and an upper non-marine sequence of alternating beds of sandstone and clay, with carbonaceous clay and lignite seams towards the bottom (Warkalli Formation). These beds are horizontally disposed and lateritised at the top. The midland portion representing the Tertiary sedimentary terrain and the western part of the Archaean terrain is extensively lateritised and the laterite is 5 to 10m thick. The coastal plain is covered by Quarternary (*Figure 1*). The geology of the district given above may be read with the “Geology of Kerala” which is given as Annexure 1 for better understanding of geological succession and stratigraphic sequence.




Figure 1: Geology and mineral resources of Kollam. (Source: District Resource map, Kollam district, Geological Survey of India)

## 6 Geomorphology

Physiographically, the district can be divided into three distinct units from west to east viz. the coastal plains, the midlands and the eastern highland regions. The coastal plains with an elevation ranging between 0-6 m amsl occur as narrow belt of alluvial deposits parallel to the coast. It has a maximum width of 90kms in the north and gradually narrows down to less than 0.5kms towards south. It is a nearly level to very gently sloping terrain depicting depositional landforms like strandlines (palaeo-beach ridges), flood plain and tidal flats. The coastal plain has a number of back waters known as kayals in Kerala – the prominent being the Ashtamudi kayal, Paravur kayal, Panmana kayal and the Sasthamkotta kayal. Among these the last one is a fresh water lagoon, while the others are brackish. To the east of coastal belt is the midland region having a rolling topography with elevations ranging from 20m to around 300m. The midland area is characterised by rugged topography formed by gently to moderately sloping spurs, moderately to steeply sloping ridges, flat and domal hills with intervening narrow valleys and broad valley floors. The midland regions show a general slope towards the western coast. To its east is the high land region. The hills have steep slopes and narrow and small summits. Highest peaks along the eastern boundary are 1200 to 1500m high. Major parts of the catchment of river Kallada and Ithikara fall within this unit. This unit occupies the maximum area of the district. The Western Ghat fringes is bounded by 300 to 600 m contours. The highest elevation is noticed at Karimalai (1758 m amsl)

## 7 Soil types

There are five major soil types encountered in the district. They are Lateritic soils, Brown Hydromorphic soils, Greyish Onattukara soils, Riverine and Coastal Alluvium and Forest Loam. Lateritic soil is the most predominant soil type of the district and it occurs in the midland and hilly areas and it is derived from laterites. Brown hydromorphic soil is confined to the valleys between undulating topography in the midlands and in the low lying areas of the coastal strip. They have been formed as a result of transportation and sedimentation of materials from adjoining hill slopes. The alluvial soil is seen in the western coastal tract of the district. The coastal alluvium is characterized by secondary soils which are sandy and sterile with poor water holding capacity. Riverine alluvium is seen along the river beds. The width of the zone increases towards the southern part of the district. Greyish Onattukara soils are purely marine deposits extending to the interior and are generally coarse in texture. Forest loamy soils are found in the eastern hilly areas of the district and are characterized by a surface layer rich in organic matter.

## 8 Land use




SI No.	Category	Area in hectares	% of total area
1	Geographical area	249100	100
2	Built up land	6926	3
3	Agriculture land	139928	56
	Forest	84293	34
4	Water bodies	11816	5
5	Waste land	4784	2
6	Others	447	0.18

## 9 Forest

Kollam District has a large area under forest. Pathanapuram, Anchal, Kottarakkara and Chadayamangalam are blocks having large areas of forest. The forest divisions are at Thenmala and Punalur. For all its natural gifts of wide rivers, dominant hills and forest, lakes and plains, there is the throb of a vibrant industrial activity that combines with equanimity the wonders of the modern age with the traces of a bygone era.

## 10 Groundwater scenario

Geohydrologically, the area comprises A, B and C zones with respect to groundwater potential as low to poor, moderate to low and fairly good respectively. The district is subdivided into Vamanapuram, Ayur, Itikara, Kallada, Pallikkal and achenkovil basins. Ground water occurs in the porous granular formations such as alluvium, laterite, the Tertiary sediments and weathered and decomposed crystalline rocks as well as in the fissures, joints and fractures in the fresh crystalline rocks. The aquifers in the district can be grouped into four distinct geological formations in which they occur viz alluvial aquifers, laterite aquifers, Tertiary sedimentary rock aquifers and crystalline rock aquifers



Source: Resource Atlas of Kerala (CESS, 1984)

## II. GEOMORPHOLOGY AND GEOHYDROLOGY

### GEOMORPHOLOGY

- 1** Nearly level to very gently sloping coastal plain
- 2** Gently to moderately sloping spurs
- 3** Moderately to steeply sloping ridges
- 4** Steep to very steep hill ranges
- 5** Valleys - gently sloping to flat bottom

### GEOHYDROLOGY

- A** Ground water potentiality low to poor
- B** Ground water potentiality moderate to low
- C** Ground water potentiality fairly good

----- Basin boundary

- I. Vamanapuram Basin
- II. Ayur Basin
- III. Ittikara Basin
- IV. Kallada Basin
- V. Pallikkal Basin
- VI. Achenkovil Basin

Figure 2: Geomorphology and geohydrology of Kollam. (Source: District Resource map, Kollam district, Geological Survey of India)




Figure 3: Land use map of Kollam district (Source: District Resource map, Kollam district, Geological Survey of India)




Figure 4: Geotechnical characteristics and natural hazards map of Kollam. (Source: District Resource map, Kollam district, Geological Survey of India)







Figure 5. Soil map of Kollam district



Source : Department of Forests, Govt. of Kerala, 2010

## 11 Natural hazards

The area comes under zone III and indicates moderate seismicity (*Figure 4*).

## 12 Mineral Resources

### 12.1 Major minerals

The district is immensely rich in mineral resources. The important minerals occurring in the district are bauxite, clay, chrysoberyl, graphite, heavy sand, mica and lime shell. The beach along Chavara-Neendakara is rich in ilmenite, rutile, zircon, monazite and garnet which offer scope of exploitation for industrial purpose. Good quality bauxite occurs in Churanad, Vadakkemuri, Adichanallur and Chittavattom areas. The  $Al_2O_3$  content varies between 40 to 50%. China clay, both of primary and secondary origin, suitable for ceramics, paper coating, textile and rubber industries, occurs in Kundara, Mulavana, Vellichikala, Perumbuzha, Kalluvathukkal and Chattannur. Gem quality chrysoberyl is reported from pegmatite veins and in the gravel and pebble beds around Karumbad, Venpakal, Talchira and Elampazhannur. In the vicinity of Karuppanthodu, Perumthottil and Changappara of Ashtamudi kayal, limeshell is reported. Thin bands of fossiliferous limestone are noticed within the Quilon Formation. Phlogopite mica occur near Punalur. The khondalite suite has graphite associated with it.

Indian Rare Earths Limited has 4 mining leases for beach sand minerals and Kerala Minerals and Metals Ltd. has one mining lease for beach sand minerals.

### 12.2 Minor Minerals

#### 12.2.1 Ordinary Earth

Ordinary earth is the common name used for the soils. Soil is made up of three main components – minerals that come from rocks below or nearby, organic matter which is the remains of plants and animals that use the soil, and the living organisms that reside in the soil. The proportion of each of these is important in determining the type of soil that is present. But other factors such as climate, vegetation, time, the surrounding terrain, and even human activities (eg. farming, grazing, gardening, landscaping, etc.), are also important in influencing how soil is formed and the types of soil that occur in a particular landscape. The formation of soils can be seen as a combination of the products of weathering, structural development of the soil, differentiation of that structure into horizons or layers, and lastly, of its movement or translocation. In fact, there are many ways in which soil may be

transported away from the location where it was first formed. Soils represent one of the most complex and dynamic natural systems and are one of the three major natural resources, other than air and water. Knowledge of their chemical, physical and biological properties is a prerequisite both for sustaining the productivity of the land, e.g. agriculture, and for conservation purposes. Soil is an integral part of a terrestrial ecosystem and fulfills numerous functions including the capacity to generate biomass and the filtering or buffering activities between the atmosphere and the groundwater in the biosphere. Soils have many important functions. Perhaps the best appreciated is the function to support the growth of agricultural and horticultural crops. Soil is the mainstay of agriculture and horticulture, forming as it does the medium in which growth and ultimately the yield of food producing crops occurs. Farmers and gardeners have worked with their soils over many centuries to produce increasing amounts of food to keep pace with the needs of a burgeoning world population. The soil's natural cycles go a long way in ensuring that the soil can provide an adequate physical, chemical and biological medium for crop growth. As well as being essential to agriculture, horticulture, forestry and natural and semi-natural systems, soil also plays an important role for our fauna. The soil itself contains millions of organisms, the exact nature and role of which we are still trying to determine. Undoubtedly, the soil flora and fauna play a vital role in cycles which are fundamental to the ability of the soil to support natural and semi-natural vegetation without additions of fertilizer and other support mechanisms. They breakdown plant debris, take in components from the atmosphere, aerate the soil together with many other functions that make the soil such an important medium.

#### *Classification of soils (ordinary earth) commonly found in the district*

The topo-lithosequence along with variation in rainfall, temperature and alternate wet and dry conditions particularly from the western coast to high ranges in the east and swift flowing rivers lead to the development of different types of natural vegetation and soil. The soils can be broadly grouped into coastal alluvium, mixed alluvium, acid saline, kari, laterite, red, hill, black cotton and forest soils. Soil map given below may be referred to find out its occurrences.

#### *Coastal Alluvium*

These soils of marine origin are identified along the coastal plains and basin lands as a narrow strip. The elevation of the coastal area is generally below 5m MSL. The area has high water table and in some areas it reaches above the surface during rainy season. The soils of the coastal plains are very deep with sandy texture. The texture generally ranges from sand to loamy sand with greyish brown to reddish brown and yellowish red colour. Sand content

ranges from 80% and clay up to 15%. Even though these soils have high water table, the water holding capacity is poor due to the predominance of sand. Coconut is the major crop in the area. Cashew and other fruit trees are also grown.

#### *Mixed Alluvium*

These soils are developed from fluvial sediments of marine, lacustrine and riverine sediments or its combinations. They occur below 20m MSL in the lowland plains, basins, valleys and along the banks of major rivers. The mixed alluvium is mainly noticed close to coastal alluvium, Kuttanad and adjacent area and kole lands of Thrissur district. The soils are frequently flooded and submerged. The soils of depressions and broad valleys are subject to occasional flooding and stagnation. The ground water table of these soils is generally high and it reaches above the surface during rainy season. A wide variation in texture is noticed in these soils. Sandy clay loam to clay is the predominant texture. Sandy loam soils are also met with. Light grey to very dark brown is the common colour of the soil. Paddy, other annuals and seasonal crops like banana, tapioca and vegetables are grown here.

#### *Laterite soil*

Laterite and laterite soil are the weathering products of rock in which several course of weathering and mineral transformations take place. This involves removal of bases and substantial loss of combined silica of primary minerals. In laterite and laterite soils, over acidic rocks, induration and zonation are more pronounced. This induration is greater if the iron content is higher. These soils mainly occur in the midlands and part of lowlands at an elevation of 10 to 100m above MSL as a strip between the coastal belt and hilly mid-upland. The area comprises of mounds and low hills with gentle to steep slopes. Laterite soils are generally suitable for most of the dry land crops. It is mainly cultivated with coconut, arecanut, banana, tapioca, vegetables, yams, pepper, pineapple, fruit trees etc. The percentage of gravel content in the soil and reduced soil depth limits the choice of crops. In laterite outcropped area with shallow soils, only cashew can be grown with vegetables.

#### *Hill Soil*

The hill soils mostly occur above an elevation of 80m MSL. The area is hilly and has highly dissected denudational hills, elongated ridges, rocky cliffs and narrow valleys. The general slope range is above 10%. The texture of these soils generally ranges from loam to clay loam with average gravel content of 10 to 50%. In addition, stones and boulders are noticed in the subsoil. These soils have reddish brown to yellowish red/strong brown colour. Generally, increase in clay content is noticed down the profile. The depth of the soil varies considerably from 60 to 200 cm depending on erodability of soil and past erosion. These soils are mostly

friable and subject to heavy soil erosion. The area is suitable for all dry land crops like rubber, coconut, arecanut and fruit trees based on the topography. Crops such as banana, pepper, pineapple, vegetables can be grown in foot slopes.

### *Forest Soil*

These soils are developed from crystalline rocks of Archaean age under forest cover. They occur along the eastern part of the State, generally above an elevation of 300m above MSL. The area is hilly and mountainous with steep slopes, escarpments, elongated rocky summits and narrow 'V' shaped valleys. The depth of the soil varies considerably depending on erosion and vegetative cover. The soils are generally immature due to slow weathering process. Rocky outcrops and stones are noticed on the surface. Gneissic boulders under different stages of weathering are noticed in the subsoil. The texture of the soil ranges from sandy clay loam to clay with reddish brown to very dark brown colour. Forest trees, shrubs and grasses are grown here.






Figure 5: Soils of Kerala



### *Mining of ordinary earth*

Usually ordinary earth is mined for levelling of ground for construction of buildings. Since ordinary earth is very important to mankind, it is not wise to mine ordinary earth for filling purposes alone. However, for the construction of roads and other infrastructure, ordinary earth is mined after obtaining quarrying permit from the Department of Mining and Geology. Mining and transporting ordinary earth/soil without the permission of Department of Mining and Geology is an offence. Department issues pass for transport of ordinary earth. Dealer's license is not issued for ordinary earth as it is not considered as a mineral mined for commercial purposes.

#### **12.2.2 Ordinary Clay (tile/brick clay)**

Clays and clay minerals occur under a fairly limited range of geological conditions and are produced by weathering of silicate minerals containing calcium, magnesium, sodium, or potassium reacting with carbonic acid, carbonates, and bicarbonates. These soluble products are removed by ground water, while the remaining elements, aluminium, silicon, and oxygen combine with water to produce stable clay minerals. The environment of formation include soil horizons, continental and marine sediments, geothermal fields, volcanic deposits, and weathering rock formations. Extensive alteration of rocks to clay minerals can produce relatively pure clay deposits that are of economic interest. Clay formed at the site of the parent rock is known as primary or residual clay; the one carried away or transported and deposited elsewhere is known as secondary clay. For obvious reasons, the former is purer with less impurity (5%–15%), while the latter may contain mica, quartz, and iron oxide as impurities. Geological factors such as conditions at the time of deposition and post-depositional changes have an important influence on the properties of sediment.

Buildings and utensils made of clay date back to the earliest periods of man's civilized development, and the use of clay is intimately associated with his history. Tile and brick kilns are closely associated with Kerala's culture and traditional architecture, which is continued in modern buildings as well.

In Kerala, tile/brick clay occurs in the wetlands/paddy fields in the lowlands and midlands. The clay extracted is used for a variety of purposes such as manufacture of roofing, flooring, and decorative tiles, wire cut (mechanically made) and ordinary bricks (manually made), and pottery wares. Studies carried out in clay mining areas of Kerala have proved that unprecedented increase in the development needs of the state and the subsequent increase in the resource extraction scenarios, especially that of clay mining, have led to rapid degradation

of the wetlands (paddy fields), which is significantly reflected in the declining agricultural productivity of the state. Mining of clays several meters below the prescribed levels, water draining from the unaffected paddy lands into the adjacent mine pits, and subsequent pumping of water for further mining impose severe problems on the hydrological regime, lowering the water table and creating severe water shortage problems in the mining areas. The additional expenditure incurred to meet the freshwater requirements of the people living in areas adjacent to mining sites is increasing year after year, which undermines the short-term economic benefits of resource extraction.

Tile and brick clay mining and its processing provide employment opportunities to a considerable section of the people in the midland and lowland areas of Kerala. Adding to this, thousands of labourers in the construction industry also indirectly depend on the products manufactured from these clays. Under these circumstances and also with respect to the demand incurred, complete restriction of extraction activities does not prove to be viable.

In the study report published by National Center for Earth Science Studies on the impact of clay mining, following recommendations were given with respect to tile/brick clay mining:

“It is of imminent importance to regulate random mining from the paddy fields/wetlands of Kerala by allowing only location-specific resource extraction under well-conceived guidelines. It is also crucial to limit the extraction of tile and brick clays to meet indigenous and local demand only. This is to save the prime agricultural land and also to increase the rice production in the area. The depth of mining should be demarcated so as to regulate mining with respect to the water table condition in the summer season. Also, adequate measures are to be taken to regenerate the natural ground water table using the stored water in the clay mine pits for irrigating the agricultural crops of the hinterland areas. This will enhance the net agricultural productivity of the area in addition to saturating the aquifer systems in the hinterlands. Awareness creation among the public about the adversities of clay mining and as well as the economic benefits of using clay bricks for construction purposes will serve in the protection of our wetlands/paddy fields. Recycling of building materials should also be considered in order to reduce mining of tile and brick clays. The abandoned clay mine areas left behind as fallow lands or water logged areas can be used for productive purposes such as fish farm ponds or irrigation ponds that promise some utility to the society. Also, suitable guidelines should be framed to streamline the tile and brick clay mining activities of the state on an eco-friendly basis.”

The Kerala Conservation of Paddy Land and Wetland Act, 2008 and Rules made thereunder which was enacted for conservation of paddy land and wetlands of Kerala imposes restrictions

in mining of tile/brick clays in such areas. The said Act and Rules are implemented by Revenue Department. In addition, Government have setup District Expert Committee to monitor and control the mining activities of ordinary clay. In Kerala Minor Mineral Concession Rules 2015, it is mandated that No Objection Certificate from the District Collector concerned, based on the recommendation of the District Expert Committee constituted by the Government in this regard, is to be produced by the applicant in the case of application for extraction of ordinary clay. In addition, Bank guarantee from any Nationalized or Scheduled Bank at the rate of Rs. 300/- (Rupees three hundred only) per cubic metre for the purpose of reclamation of pits that will be formed after quarrying in the area permitted, in respect of application for extraction of ordinary clay. Based on the request of the entrepreneurs working in tile/brick clay based industry, Government have instructed the Department of Mining and Geology to carry out survey to identify the mineable tile/brick clay deposits of Kerala and the work in this respect is progressing.

### **12.2.3 Ordinary Sand**

In Kerala Minor Mineral Concession Rules, 2015, the ordinary sand is defined as sand used for non-industrial purpose. This includes both river sand and sand excavated from inland areas like palaeo-channels. Since a separate Act has been enacted by Government of Kerala namely, The Kerala Protection of River Banks and Regulation of Removal of Sand Act, 2001 (hereafter referred to as Sand Act, 2001) and since the mining of river sand is controlled by Revenue Department by virtue of the powers conferred by the said Act and the Rules made thereunder, the Department of Mining and Geology now regulates the mining of sand which do not comes under the purview of Sand Act, 2001.

The ordinary sand (other than river sand) occurs in the palaeo-channels. The word palaeo-channel is formed from the words “palaeo” or “old,” and channel; i.e., a palaeo-channel is an old channel. Palaeo-channels are deposits of unconsolidated sediments or semi-consolidated sedimentary rocks deposited in ancient, currently inactive river and stream channel systems. These are typical riverine geomorphic features in a location representing drainage streams, rivers, rivulets which were flowing either ephemeral or perennial during the past time and now stands either buried or lost or shifted due to tectonic, geomorphologic, anthropogenic process/activities, as well as climatic changes. When a channel ceases to be part of an active river system, it becomes a palaeo-channel. In order to tap the ordinary sand occurring in palaeo-channels, the Department entrusted the study of identification of palaeo-channels in major river basins of Kerala to Geological Survey of India (GSI). GSI resorted to remote

sensing studies using satellite imageries and delineated some of the palaeo-channels. However, since such deposits falls in paddy land/wetlands of Kerala, it is difficult to extract such sand on account of restrictions imposed by various Acts and Rules.

The Kerala Conservation of Paddy Land and Wetland Act, 2008 and Rules made thereunder which was enacted for conservation of paddy land and wetlands of Kerala imposes restrictions in mining of ordinary sands occurring in wetlands and paddy fields. The said Act and Rules are implemented by Revenue Department. In addition, Government have setup District Expert Committee to monitor and control the mining activities of ordinary sand. In Kerala Minor Mineral Concession Rules 2015, it is mandated that No Objection Certificate from the District Collector concerned, based on the recommendation of the District Expert Committee constituted by the Government in this regard, is to be produced by the applicant in the case of application for extraction of ordinary sand. In addition, Bank guarantee from any Nationalized or Scheduled Bank at the rate of Rs. 300 (Rupees three hundred only) per cubic metre for the purpose of reclamation of pits that will be formed after quarrying in the area permitted, in respect of application for extraction of ordinary sand.

The mining of ordinary sand from palaeo-channels also case some environmental concerns. Since sand is a good aquifer, the mining of aquifer system poses threat to ground water availability in surrounding areas. However in certain cases, the mining of such sand from paddy lands increase the productivity of paddy as excess sand in the paddy lands are not good for paddy.

In Kerala, due to shortage of river sand and ordinary sand occurring in palaeo-channels, the construction industry now uses manufactured sand obtained by crushing of crystalline rocks.

**It may be noted that since the Revenue Department is taking care of all types of mining activities related to river sand and since sand auditing and other studies are carried out under the aegis of the Revenue Department, this report shall not be used for the purpose of obtaining prior environmental clearance for mining of river sand.**

#### **12.2.4 Laterite**

Laterite is a soil and rock type rich in iron and aluminium, and is commonly considered to have formed in hot and wet tropical areas. Nearly all laterites are of rusty-red coloration, because of high iron oxide content. They develop by intensive and long-lasting weathering of the underlying parent rock. Tropical weathering is a prolonged process of chemical weathering which produces a wide variety in the thickness, grade, chemistry and ore mineralogy of the

resulting soils. The majority of the land area containing laterites is between the tropics of Cancer and Capricorn.

Angadipuram Laterite is a National Geological Monument identified in Angadipuram town in Malappuram district. The special significance of Angadipuram to laterites is that it was here that Dr. Francis Buchanan-Hamilton, a professional surgeon, gave the first account of this rock type, in his report of 1807, as "indurated clay", ideally suited for building construction. This formation falls outside the general classification of rocks namely, the igneous, metamorphic, or sedimentary rocks but is an exclusively "sedimentary residual product". It has a generally pitted and porous appearance. The name laterite was first coined in India, by Buchanan and its etymology is traced to the Latin word "letritis" that means bricks. This exceptional formation is found above parent rock types of various composition namely, charnockite, leptynite, anorthosite and gabbro in Kerala. The laterite profiles in different types of rocks vary depending on the composition of parent rock. For example in Charnockites, the thickness of the profile ranges from 2 m to 10 m with humus zone on the top with thin pebbly zone (with ferruginous pellets in clayey matrix), underlain by vermicular laterite with tubular cavities of various shapes and size filled with kaolinitic clay. This is followed by thin layer of lithomarge. Further below completely weathered, partly weathered or fresh parent rock occur. In some places one can see hard duricrust at the top.

The mineralogical study of laterites reveals that all the silicate minerals have been transformed to a mixture of goethite, hematite and kaolinite in laterite samples developed over charnockite. Further studies revealed that pyroxenes have been altered to goethite while feldspars gave rise to kaolinite. Quartz is cracked, eroded and disintegrated. Monazite and Zircons are found as accessory minerals.

Laterite and bauxite show a tendency to occur together. Aluminous laterites and ferruginous bauxites are quite common. The most common impurity in both is silica. Laterite gradually passes into bauxite with decrease in iron oxide and increase in aluminium oxide. The laterite deposits may be described on the basis of the dominant extractable minerals in it: (i) aluminous laterite (bauxite), (ii) ferruginous laterite (iron ore), (iii) manganiferous laterite (manganese ore), (iv) nickeliferous laterite (nickel ore) and (v) chromiferous laterite (chrome ore). Laterite with  $Fe_2O_3:Al_2O_3$  ratio more than one, and  $SiO_2:Fe_2O_3$  ratio less than 1.33 is termed as ferruginous laterite, while that having  $Fe_2O_3:Al_2O_3$  ratio less than one and  $SiO_2:Al_2O_3$  ratio less than 1.33 is termed as aluminous laterite. Laterite can be considered as poly-metallic ore as it is not only the essential repository for aluminium, but also a source of iron, manganese, nickel and chromium. Furthermore, it is the home for several trace elements like gallium and

vanadium which can be extracted as by-products. In Kerala laterites are extracted as building stones which are used for construction of building. Laterite as a building stone possesses one advantage that it is soft when quarried and can be easily cut and dressed into blocks and bricks which on exposure to air become hard. In addition, laterite (aluminous laterite) is extracted for industrial purposes (for eg. Cement industry). In addition to aluminous laterite, bauxites are also mined in Kerala. Hence, while granting mineral concession for laterite it is necessary to carry out the chemical analysis to establish whether the mineral is bauxite or aluminous laterite.

#### **12.2.5 Granite Dimension Stone and Granite (building stone)**

For administrative purpose the hard crystalline rocks which do not have any economic minerals are classified as granite dimension stones and granite (building stones). The definition given in the Kerala Minor Mineral Concession Rules 2015 is as follows:-

‘Granite dimension stones include all types of granites, dolerite, charnockite, leptynite and other crystalline rocks of Acid, Intermediate, basic and ultra basic groups of igneous and metamorphic origin which are suitable for cutting to pre-determined sizes, polishing, carving and amenable for making value-added products in decorative monumental and ornamental fields of industry as a high-value item. Granite (building stone) include all those group of rocks specified above which are not suitable for using as dimension stones as specified therein, but can be used as ordinary building stones, road metal, rubble and ballasts after breaking into irregular pieces by blasting or otherwise as low value item. The Rules insists that the rocks having the quality of granite dimension stone shall not be quarried for granite building stone as these two types of rocks have different values/royalties’.

The major granite dimension stone occurrence in the district forms part of Charnockite-Khondalite belt and has colour ranging from pale green with mottled red, bluish green with cordierite, deep dark green, greyish white.

All Archaean and Proterozoic rocks of Kerala (refer section on Geology of Kerala) which are not listed above as granite dimension stone falls under the category of granite (building stone) and are found below ordinary earth/laterites/and other sedimentary rocks. In some cases such rocks are exposed as hillocks without any overburden.

### **13 Details of minor mineral concessions and revenue collection**

Permission for mining will be granted on case to case basis on ascertaining the availability at the site and only if conditions stipulated in the KMMC Rules 2015 are satisfied (The reader may refer the KMMC Rules 2015 available in the website [www.dmg.kerala.gov.in](http://www.dmg.kerala.gov.in) for more details in this regard). The concession will be granted only if other statutory licenses like Environmental Clearance, Explosive Licence, consent to operate issued by State Pollution Control Board, NOC issued by Revenue Department (as the case may be), Dangerous and Offensive Trade Licence issued by Local Self Government Institutions, NOC related to Coastal Regulation Zone (as the case may be), NOC issued by Forest (as the case may be) etc. The mineral concession will not be granted in the ecologically sensitive areas, ecological fragile zones etc. The details of valid minor mineral concessions issued by the Department are given in the table below:-



**Table1: Details of revenue collection for the period 2013-'14, 2014-'15 and 2015-'16**

**Minor Minerals (2013-'14)**

No.	Particulars	Granite Building Stone	Laterite Building Stone	Brick Clay	Ordinary Sand	River Sand	Ordinary Earth	Total
1	Royalty	56000	36000	40000	55000		5892273	6079273
2	Arrear Royalty	3456933	326102	30920	8480		196180	4018615
3	Surface Rent	2800	5400	40				8240
4	Fee for Quarrying Permit	5600	11600	400			86000	103600
5	Fee for Dealer's Licence	1210000						1210000
6	Consolidated Royalty	3515000	719800					4234800
7	Other Departments					893376		893376
8	RMCU Compounding	650000						650000
<b>Total</b>		<b>8896333</b>	<b>1098902</b>	<b>71360</b>	<b>63480</b>	<b>893376</b>	<b>6174453</b>	<b>17197904</b>

### Minor Minerals (2014-'15)

No.	Particulars	Granite Building Stone	Laterite Building Stone	Brick Clay	Ordinary Sand	River Sand	Ordinary Earth	Total
1	Royalty	48000		102112			2054690	2204802
2	Arrear Royalty	6008871	145291	7400	43360		415860	6620782
3	Surface Rent	14300	4400					18700
4	Fee for Quarrying Permit	28400	8800	200			3200	40600
5	Fee for Dealer's Licence	2001000						2001000
6	Consolidated Royalty	14505000	750000					15255000
7	Other Departments					249789		249789
8	RMCU Compounding	2254000						2254000
<b>Total</b>		<b>24859571</b>	<b>908491</b>	<b>109712</b>	<b>43360</b>	<b>249789</b>	<b>2473750</b>	<b>28644673</b>

### Minor Minerals (2015-'16)

No.	Particulars	Granite Building Stone	Laterite Building Stone	Ordinary Earth	Brick Clay	River sand
1	Royalty	26217760	321525	7424470	287440	280000
2	Arrear Royalty	5225546	473985	495200	100900	
3	Surface Rent	30800	2800			
4	Registration fee for D.L	54000				
6	D.L.Fee	3348000				
7	Application Fee for Quarrying Permit	95000	33000			
10	Fee for Dealer's Licence	36000				
11	Consolidated Royalty	24038690	3935571			
15	RMCU Compounding	2300000				
<b>Total</b>		<b>61345796</b>	<b>4766881</b>	<b>7919670</b>	<b>388340</b>	<b>280000</b>

**List of valid mineral concessions**

<b>Table 2a: List of Quarrying Lease granted for Granite building stone</b>										
<b>Sl. No.</b>	<b>Concession holder's name and address</b>	<b>Concession no.</b>	<b>Mineral</b>	<b>Survey no</b>	<b>Village</b>	<b>Taluk</b>	<b>District</b>	<b>Area (ha)</b>	<b>valid from</b>	<b>valid to</b>
1	M.K. Biju, Karthika, Kuttikkad, Kadackal.	752/2005-06/ 2264/M3/06	Granite building stone	432/4	Kummil	Kottarakkara	Kollam	1.1820 Hect	22-03-2006	21-03-2018
2	M.K. Biju, Karthika, Kuttikkad, Kadackal.	751/2005-06/ 1684/M3/06	Granite building stone	161	Ittiva,	Kottarakkara	Kollam	0.850 Hect.	22-03-2006	22-03-2018
3	M.K. Biju, Karthika, Kuttikkad, Kadackal.	750/2005-06/ 2266/M3/06	Granite building stone	158/6, 12-2, 158/5	Ittiva,	Kottarakkara	Kollam	0.7277 hect.	22-03-2006	21-03-2018

4	Gopalakrishnakurup, G.K. Industries, Kalluvathukkal, Kollam	562/2005- 06/ 9571/M3/05	Granite building stone	359/2, 359/3-1, 359/15	Kalluvathukkal	Kollam	Kollam	0.7265 hect.	07-01-2006	06-01- 2018
5	M. Abraham, Charuvila Puthen Veedu, Ayoor	806/2010- 11/ 1687/M3/11	Granite building stone	151/4-2, 4-3, 152/1-2, 1-3	Valakom	Kottarakkara	Kollam	2.0566 hect.	08-03-2011	06-03- 2017
6	Sri. Saji M. Mathew, Muthilakkattil House, Aryankavu P.O., Kollam	616/2009- 10/ 5595/M3/09	Granite building stone	577/1 pt.,	Aryankavu	Pathanapuram	Kollam	48.56 Ares	04-03-2010	09-03- 2022
7	M/s. Poabs Entrprises (P) Ltd, Kuttoor P.O., Thiruvalla, Pathanathitta	663/2010- 11/ 9943/M3/10	Granite building stone	270/2, 269/2, 272/2-2, 273/3, 274/1-2	Chithara	Kottarakkara	Kollam	2.1649 hect.	03-01-2011	27-01- 2017

8	M/s. Poabs Entrprises (P) Ltd, Kuttoor P.O., Thiruvalla, Pathanathitta	940/2011-12/684/M3/12	Granite building stone	266/15,24, 16-1, 16-2, 271/7-1, 7-2, 7-3, 272/5	Chithara	Kottarakkara	Kollam	1.0397 hect.	28/3/2012	04-01-2017
9	M/s. Poabs Entrprises (P) Ltd, Kuttoor P.O., Thiruvalla, Pathanathitta	928/2011-12/685/M3/12	Granite building stone	267/5, 268/2-1, 2-2, 3-2-1, 3-2-2, 5-2, 6,7,9-2, 8-1, 8-2, 4, 4-2, 4-3	Chithara	Kottarakkara	Kollam	2.4703 hect.	24/3/2012	04-01-2019

<b>Sl. No</b>	<b>Concession holder's name and address</b>	<b>Concession no.</b>	<b>Survey no</b>	<b>Village</b>	<b>Taluk</b>	<b>Area (ha)</b>	<b>valid from</b>	<b>valid to</b>	<b>Consolidated royalty (Rs.)</b>
1	G.Rajeevan, Vazhappally Padinjattethil, Mannadi.P.O, Kottarakkara	64/2015-16/MM/GBS/CRPS/DOQ/S2/1321/2015	501/1-7, 501/4-2, 501/5-3, 501/5-4	Kareepra	Kottarakkara	27.93Ares	7/12/15	6/12/16	300000
2	Shajahan, Amina Manzil, Nedumankavu.P.O	66/2015-16/MM/GBS/CRPS/DOQ/S2/2701/2015	383/1-4, 383/1-5	Valakom	Kottarakkara	28.33 Ares	28/12/15	27/12/16	300000
3	Aliyarukutty, M.S. House, Vakkanadu, Kareepra.	76/2015-16/MM/GBS/CRPS/DOQ/S2/2657/2015	9/8-1, 8-2	Veliyam	Kottarakkara	46.12 Ares	16/1/2016	6/12/2016	700000
4	Jose Abraham, Panamthottathilveedu, Kottara, Meeyannoor.P.O	79/2015-16/MM/GBS/CRPS/DOQ/S2/2806/2015	364/17, 364/5pt, 364/6-2, 364/8pt	Veliyam	Kottarakkara	14.53Ares	19/1/16	16/12/16	150000
5	Ambujakshan, Kavuvilaveedu, Karikkal, Karimpinuzha.P.O, Puthoor	80/2015-16/MM/GBS/CRPS/DOQ/S2/158/2016	465/1-4	Neduvathoor	Kottarakkara	5.00 Ares	02-02-2016	16/12/16	75000
6	Balan, Pulivilaputhenveedu, Vendar.P.O, Puthoor	81/2015-16/MM/GBS/CRPS/DOQ/S2/1692/2015	372/1, 373/6, 373/6-3	Puthoor	Kottarakkara	50.00 Ares	02-02-2016	16/12/16	700000
7	V.Rajan, Puthuvlputhenveedu, Poovattoor East, Kalayapuram	83/2015-16/MM/GBS/CRPS/DOQ/S2/2777/2015	201/2-3	Puthoor	Kottarakkara	25.24 Ares	02-03-2016	16/12/16	300000



8	G.Raju, Gayathribhavan, Mylamkulam, Puthoor	85/2015- 16/MM/GBS/CRPS/ DOQ/S2/2231/2015	216/8	Puthoor	Kottarakkara	21,58 Ares	02-03-2016	16/12/16	300000
9	Saleem.M, Saleem Manzil, Nileswaram.P.O, Neduvathoor	86/2015- 16/MM/GBS/CRPS/ DOQ/S2/2881/2015	241/12, 11-2	Neduvathoor	Kottarakkara	30.56 Ares	02-03-2016	16/12/16	500000
10	Sahadevan, Salraj Bhavan, Vendar.P.O	87/2015- 16/MM/GBS/CRPS/ DOQ/S2/2807/2015	458/7, 450/4	Neduvathoor	Kottarakkara	28.75 Ares	20/2/2016	16/12/16	300000
11	Binu.S, Bineeshbhavan, Nedumankavu.P.O	88/2015- 16/MM/GBS/CRPS/ DOQ/S2/2987/2015	501/5-2, 5-5	Kareepra	Kottarakkara	09.58 Ares	26/2/16	16/12/16	75000
12	S.Shaji, Thekkuvila Melathil, Kudavattoor	90/2015- 16/MM/GBS/CRPS/ DOQ/S2/354/2015	16/3-2, 12/9-2	Veliyam	Kottarakkara	14.63 Ares	29/2/16	16/12/16	150000
13	Hari.N, Mukaluvilathekkathil, Ulakode, Vakkanadu	91/2015- 16/MM/GBS/CRPS/ DOQ/S2/353/2015	49/2-2pt, 49/2-3, 49/4	Veliyam	Kottarakkara	17.31 Ares	29/2/16	16/12/16	150000
14	Raghunathan, Rahul Nivas, Paruthiyara, Odanavattom	92/2015- 16/MM/GBS/CRPS/ DOQ/S2/1363/2015	3/3	Veliyam	Kottarakkara	44.44 Ares	03-01-2016	16/12/16	700000
15	Seeba.T.S, Plankala veedu, Edakkidam.P.O	94/2015- 16/MM/GBS/CRPS/ DOQ/S2/259/2016	396/7	Ezhukone	Kottarakkara	13.58 Ares	03-03-2016	6/12/16	150000

**Table 2c: List of Quarrying Permit granted under CRPS for Granite Building Stone (2016-'17)**

Sl. No	Concession holder's name and address	Concession no.	Survey no	Village	Taluk	Area (ha)	valid from	valid to	Consolidated royalty (Rs.)
1	Chandranandan, Kavanadupalavila veedu, Vendar	2/2016- 17/MM/GBS/CRPS/DOQ/S2/6 48/2016	334/3, 5-2, 4,5,7	Puthoor	Kottarakkara	32.61 Ares	13/4/16	6/12/16	500000
2	Chandranandan, Kavanadupalavila veedu, Vendar	5/2016- 17/MM/GBS/CRPS/DOQ/S2/1 793/2016	427/6,2,7,9, 4	Neduvathoor	Kottarakkara	45.00 Ares	06-06- 2016	6/12/16	700000
3	Jaleel, Punnavilaveedu, Vakanad, Kareepra	6/2016- 17/MM/GBS/CRPS/DOQ/S2/5 90/2016	9/12	Veliyam	Kottarakkara	17.70Ares	13/6/2016	6/12/16	150000
4	G.Rajeevan, Vazhappallipadinjattethil, Mannadi.P.O	8/2016- 17/MM/GBS/CRPS/DOQ/S2/1 219/2016	4/7, 5/8-2, 8-3, 8-8, 8- 9	Veliyam	Kottarakkara	25.21 Ares	15/6/2016	6/12/16	300000
5	Udayan.S, Navami, Kayila	10/2016- 17/MM/GBS/CRPS/DOQ/S2/1 189/2016	465/20	Veliyam	Kottarakkara	111.33 Ares	20/6/2016	6/12/16	150000
6	Udayan.S, Navami, Kayila	10/2016- 17/MM/GBS/CRPS/DOQ/S2/1 189/2016	465/20	Veliyam	Kottarakkara	11.33 Ares	20/6/2016	6/12/16	150000
7	Omanakuttan, Muralisadanam, Vakkanadu.P.O	11/2016- 17/MM/GBS/CRPS/DOQ/S2/5 35/2016	404/1	Ezhukone	Kottarakkara	17.00 Ares	20/6/2016	6/12/16	150000

8	Najeem, Punnabila Vadakkathil, Vakkanadu.P.O, Kareepra	17/2016- 17/MM/GBS/CRPS/DOQ/S2/1 668/2016	10/2	Veliyam	Kottarakkara	22.96Ares	28/6/2016	6/12/16	300000
9	N.Asokan, Ajeeshbhavan, Perayam, Umayanalloor	18/2016- 17/MM/GBS/CRPS/DOQ/S2/1 540/2016	9/1	Veliyam	Kottarakkara	12.50Ares	29/6/2016	6/12/16	150000
10	Rajeev, Rajeevam, Velichikala, Adichanalloor	19/2016- 17/MM/GBS/CRPS/DOQ/S2/2 538/2016	151/1-1	Valakom	Kottarakkara	19.54Ares	30/6/2016	6/12/16	150000
11	Remeshkumar, Aravindavilasam, Panthaplavu, Pattazhy	20/2016- 17/MM/GBS/CRPS/DOQ/S2/7 38/2016	113/3-2	Kalayapuram	Kottarakkara	8.09 Ares	07-02- 2016	1/7/2017	75000
12	Sureshkumar, Erattakkalayil, Manjalloor.P.O, Pathanapuram	22/2016- 17/MM/GBS/CRPS/DOQ/S2/5 04/2016	441/5	Mylam	Kottarakkara	10.45Ares	07-04- 2016	3/7/2017	150000
13	R.Santhosh, SanthoshNivas, Chirakkathazham.P.O	23/2016- 17/MM/GBS/CRPS/DOQ/S2/1 776/2016	3/9	Veliyam	Kottarakkara	13.40Ares	07-08- 2016	7/7/2016	150000
14	Nadarajan, Pazhavara Veedu, Kayila	29/2016- 17/MM/GBS/CRPS/DOQ/S2/1 704/2016	469/3, 3-2	Veliyam	Kottarakkara	25.60	30/7/16	29/7/2017	300000
15	Sundaresan, Padmadeepam, Mylamkulam	30/2016- 17/MM/GBS/CRPS/DOQ/S2/1 850/2016	201/4	Puthoor	Kottarakkara	19.43	30/7/16	29/7/2017	150000

16	Sundaresan, Padmadeepam, Mylamkulam	31/2016- 17/MM/GBS/CRPS/DOQ/S2/1 849/2016	201/6	Puthoor	Kottarakkara	20.30	30/7/16	29/7/2017	300000
17	Shaji, Thekkuvilamelathil, Kudavttoor	32/2016- 17/MM/GBS/CRPS/DOQ/S2/1 931/2016	12/8-7-2, 12/8-5, 12/8-4, 12/8-7, 12/8-6	Veliyam	Kottarakkara	18.78	30/7/16	29/7/2017	150000
18	Shaji, Thekkuvilamelathil, Kudavttoor	33/2016- 17/MM/GBS/CRPS/DOQ/S2/1 932/2016	9/5-3, 9/5-2	Veliyam	Kottarakkara	12.15	30/7/16	29/7/2017	150000
19	Jayakumar, Melevila Veedu, Kottara, Meyannoor.P.O, Kollam	34/2016- 17/MM/GBS/CRPS/DOQ/S2/1 758/2016	1/5-1	Veliyam	Kottarakkara	30.42 Ares	08-06- 2016	5/8/2017	500000
20	G.Raju Gayathribhavan Mylamkulam, Puthoor	35/2016- 17/MM/GBS/CRPS/DOQ/S2/1 933/2016	110/6, 110/2	Kalayapuram	Kottarakkara	14.20 Ares	08-08- 2016	7/8/2017	150000
21	Rajeevan, VazhappallyPadinjattethil , Mannadi	36/2016- 17/MM/GBS/CRPS/DOQ/S2/3 02/2016	151/1-2	Valakom	Kottarakkara	30.00 Ares	08-08- 2016	7/8/2017	300000
22	Najeem, Punnavilavadakkathil, Vakkanadu.P.O	37/2016- 17/MM/GBS/CRPS/DOQ/S2/2 093/2016	13/14, 13/17	Veliyam	Kottarakkara	33.27 Ares	08-10- 2016	9/8/2017	500000
23	JoseDaniel, Poikavilaveedu, Vengoor	38/2016- 17/MM/GBS/CRPS/DOQ/S2/3 57/2016	381/5-2	Valakom	Kottarakkara	12.95 Ares	08-11- 2016	10/8/2017	150000

24	Suresh.G, Sindhuvilasam, Kottathala	39/2016-17/MM/GBS/CRPS/DOQ/S2/202/2016	31/1	Puthoor	Kottarakkara	21.96 Ares	26/8/2016	25/8/2017	300000
25	G.Rajeevan, Vazhappallypadinjattethil, Mannadi	41/2016-17/MM/GBS/CRPS/DOQ/S2/1842/2016	202/1	Puthoor	Kottarakkara	19.05 Ares	19/9/2016	18/9/2017	150000
26	Nadarajan, Pazhavara veedu, Kayila, Veliyam	42/2016-17/MM/GBS/CRPS/DOQ/S2/2027/2016	469/1, 469/6, 469/7	Veliyam	Kottarakkara	2840 Ares	28/9/2016	27/9/2017	300000
27	Najeem, Punnavilavadakkathil, Vakkanadu.P.O, Kareepra	43/2016-17/MM/GBS/CRPS/DOQ/S2/2092/2016	10/4	Veliyam	Kottarakkara	29.41 Ares	30/9/2016	29/9/2017	300000
28	Thulaseedharan Pillai, Aswathibhavanam, Vllam, Kottarakkara.P.O, Kottarakkara	44/2016-17/MM/GBS/CRPS/DOQ/S2//2016	498/10-1	Neduvathoor	Kottarakkara	31.10 Ares	30/9/2016	29/9/2017	500000
29	Suresh.G, Sindhuvilasam, Kottathala.P.O	45/2016-17/MM/GBS/CRPS/DOQ/S2/201/2016	27/7	Puthoor	Kottarakkara	16.00 Ares	10-07-2016	06/10/2017	150000
30	Sundaresan, Padmadeepam, Mylamkulam, Puthoor	46/2016-17/MM/GBS/CRPS/DOQ/S2/2433/2016	114/2-1	Puthoor	Kottarakkara	27.50 Ares	14/10/2016	13/10/2017	300000
31	Gopalakrishnan, Kottoor House, Nedumpana.P.O, Kollam	47/2016-17/MM/GBS/CRPS/DOQ/S2/2587/2016	9/10	Veliyam	Kottarakkara	7.10 Ares	18/10/2016	17/10/2017	75000

32	Aliyarukutty, M.S.House, Vakkanadu.P.O, Kareepra	48/2016- 17/MM/GBS/CRPS/DOQ/S2/2 095/2016	9/3, 9/3/2, 9/3/3	Veliyam	Kottarakkara	42.00 Ares	19/10/201 6	18/10/201 7	700000
33	N.Asokan, Murahara, Mukhathala.P.O, Kollam	49/2016- 17/MM/GBS/CRPS/DOQ/S2/2 784/2016	<b>152/5,</b> <b>152/2</b>	Valakom	Kottarakkara	49.82 Ares	26/10/201 6	25/10/201 7	700000
34	Ravikumar, Muraleesadanam, Elayam, Vakkanadu.P.O	50/2016- 17/MM/GBS/CRPS/DOQ/S2/2 708/16	226/8-2, 226/8-3, 226/8-7	Veliyam	Kottarakkara	33.5 Ares	14/11/201 6	13/11/201 7	500000



**Table 2d: List of short term permits granted for Granite Building Stone (2016-'17)**

Sl. No.	Concession holder's name and address	Concession no.	Survey no.	Village	Taluk	District	Area (ha)	valid from	valid to	Consolidated royalty (Rs.)
1	Rajeevan, Vazhappallil House, Mannadi	12/2016-17/MM/GBS/STP/DO Q/S2/245/2016	379/10, 13-3, 14, 14-2, 17, 19, 383/1	Valakom	Kottarakkara	Kollam	76.30 Ares	24/6/2016	12-06-2016	600000
2	M.Sajeev, Roadvila Veedu, Ottumala, Kattadi.P.O	15/2016-17/MM/GBS/STP/DO Q/S2/2818/2016	225/20-4, 225/6-2, 225/8	Velinalloor	Kottarakkara	Kollam	89.79 Ares	13/7/2016	07-12-2017	6,00,000
3	DillySukhy, Anandapuri Blue Metals, Moonnumukku.P.O, Chithara	16/2016-17/MM/GBS/STP/DO Q/S2/1765/2016	111/8-2, 113/1-2, 113/1-3	Chithara	Kottarakkara	Kollam	62.45	22/7/2016	21/7/2017	1612800
4	Dileeshkumar, Chaprayil, Maranadu.P.O,	18/2016-17/MM/GBS/STP/DO Q/S2/1935/2016	240/7, 240/4-1	Ezhukone	Kottarakkara	Kollam	99.00 Ares	26/8/2016	25/7/2017	900000
5	Anilkumar, Ambalakkara, Pulamon.P.O, Kottarakkara	19/2016-17/MM/GBS/STP/DO Q/S2/1851/2016	216/1-2	Pooyappally	Kottarakkara	Kollam	93.37 Ares	29/8/2016	28/7/2017	921600

6	Tinson John, Aiswarya Granites, Arkkannoor.P.O, Elavinmoodu	20/2016- 17/MM/GBS/STP/DO Q/S2/2338/16	265/1, 265/1- 2pt	Elamadu	Kottarakkara	Kollam	66.83Ares	09-06- 2016	08-05- 2017	900000
7	Tinson John, Aiswarya Granites, Arkkannoor.P.O, Elavinmoodu	21/2016- 17/MM/GBS/STP/DO Q/S2/2341/16	117/5-2, 117/2, 117/3	Elamadu	Kottarakkara	Kollam	70.75 Ares	09-06- 2016	08-05- 2017	900000
8	Tinson John, Aiswarya Granites, Arkkannoor.P.O, Elavinmoodu	22/2016- 17/MM/GBS/STP/DO Q/S2/2342/16	262/1	Elamadu	Kottarakkara	Kollam	98.00 Ares	09-06- 2016	08-05- 2017	900000
9	Tinson John, Aiswarya Granites, Arkkannoor.P.O, Elavinmoodu	23/2016- 17/MM/GBS/STP/DO Q/S2/2342/16	264/2-2, 264/5-2	Elamadu	Kottarakkara	Kollam	99.52 Ares	09-06- 2016	08-05- 2017	900000
10	S.A.Rasheed, Paravila Veedu, Cheriyavelinalloor	24/2016- 17/MM/GBS/STP/DO Q/S2/2328/16	151/3	Elamadu	Kottarakkara	Kollam	67.60 Ares	19/9/2016	18/3/2017	384000
11	Abdulsalam, Punnivilavadakkathil, Vakkanad.P.O, Kareepra	25/2016- 17/MM/GBS/STP/DO Q/S2/2873/16	01-Mar	Veliyam	Kottarakkara	Kollam	60.20 Ares	10-07- 2016	31/3/2017	6,00,000

**Table 2e: List of Quarrying Permit granted under CRPS for Laterite Building Stone during 2015-'16**

Sl. No	Concession holder's name and address	Concession no.	Survey no	Village	Taluk	Area (ha)	valid from	valid to	Consolidated royalty (Rs.)
1	Sudarsanan, Mukaluvilaveedu, Punakkanoor, Alummodu	27/2015- 16/MM/LTBS/CRPS/DOQ/S 2/2534/2015	328/15- 3	Kareepra	Kottarakkara	9.50 Ares	30/11/2015	29/11/16	75000
2	Sabumon, Kozhikunnath Veedu, Kottathala.P.O, Kottarakkara	28/2015- 16/MM/LTBS/CRPS/DOQ/S 2/1294/2015	30/5	Odanavattom	Kottarakkara	9.10 Ares	5/12/2015	4/12/16	75000
3	Muhammad Sibi, Kailasamangalam, Vendar.P.O, Kottathala	29/2015- 16/MM/LTBS/CRPS/DOQ/S 2/2539/2015	467/9-2	Neduvathoor	Kottarakkara	9.00 Ares	5/12/2015	4/12/16	75000

4	Madhu.R, Madhu Nivas, Northmynagappally, Karunagappally	30/2015-16/MM/LTBS/CRPS/DOQ/S 2/2246/2015	459/8	Neduvathoor	Kottarakkara	11.20 Ares	14/12/15	13/12/16	75000
5	Mohanachandran, Thadathivilaputhenvedu, Karuveli, Kottarakkara	31/2015-16/MM/LTBS/CRPS/DOQ/S 2/88/2016	222/2, 222/2-2	Pavithreswar am	Kottarakkara	9.59 Ares	16/1/16	15/1/17	75000
6	Santhosh, Santhosh nivas, Chirakkarathazham, Kollam	32/2015-16/MM/LTBS/CRPS/DOQ/S 2/2785/2015	184/9,5, 7	Chirakkara	Kollam	4.05 Ares:	16/1/16	15/1/17	75000
7	Donbosco, Chariv Poika, Kumbalam.P.O	33/2015-16/MM/LTBS/CRPS/DOQ/S 2/2347/2015	97/11	Pavithreswar am	Kottarakkara	9.90Ares	19/1/16	18/1/17	75000
8	Prasanth, Kallingal veedu, Meenadu East, Chathannoor	34/2015-16/MM/LTBS/CRPS/DOQ/S 2/1892/2015	144/18/ 5, 144/8/6	Veliyam	Kottarakkara	7.49 Ares	19/1/16	18/1/17	75000

9	Indu.P, Kuzhgvdfnd]hglldlvd] br'd, Ambalathumkala	35/2015- 16/MM/LTBS/CRPS/DOQ/S 2/2256/2015	528/2-1	Naduvathoor	Kottarakkara	8.5 Ares	19/1/16	18/1/17	75000
10	Surendran, Kanjiramvila veedu, S.N.Puram	36/2015- 16/MM/LTBS/CRPS/DOQ/S 2/322/2016	453/5	Pavithreswar am	Kottarakkara	6.07 Ares	19/2/16	18/2/17	75000
11	K.K.Rajagopal, Paurname, Meenadu East, Chathannoor	37/2015- 16/MM/LTBS/CRPS/DOQ/S 2/421/2016	559/6	Neduvathoor	Kottarakkara	3.24 Ares	26/2/16	25/2/2017	75000
12	Negarudeen, Charuvilaputhenveedu , Veliyam, Kottarakkara	38/2015- 16/MM/LTBS/CRPS/DOQ/S 2/485/2016	56/14, 56/30	Odanavattom	Kottarakkara	8.90 Ares	1/3/2016	28/2/2017	75000
13	Prasad, Mannathveedu, Mynagappally	39/2015- 16/MM/LTBS/CRPS/DOQ/S 2/2684/2016	588/4	Kunnathoor	Kunnathoor	9.82 Ares	3/3/2016	2/3/2017	75000

14	Jacob John, Panayilveedu, Odanavattom	40/2015- 16/MM/LTBS/CRPS/DOQ/S 2/2936/2016	3/1-3-2, 3/1-3	Odanavattom	Kottarakkara	19.02Ar es	8/3/2016	7/3/2016	150000
15	Johnson, Kanjirakkal Veedu, Odanavattom	41/2015- 16/MM/LTBS/CRPS/DOQ/S 2/2937/2015	3/11	Odanavattom	Kottarakkara	3.85 Ares	8/3/2016	7/3/2016	75000
16	L.Vijayan, RajeshBhavan, Karuveli.P.O, Ezhukone	42/2015- 16/MM/LTBS/CRPS/DOQ/S 2/481/2016	530/5-2	Ummannoor	Kottarakkara	9.90 Ares	17/3/2016	16/3/2017	75000

**Table 2f: List of Quarrying Permit granted under CRPS for Laterite Building Stone during 2016-'17**

Sl. No	Concession holder's name and address	Concession no.	Survey no	Village	Taluk	Area (ha)	valid from	valid to	Consolidated royalty (Rs.)
1	Deepu.G, Kanjamkunnilveedu, Edakkidam	1/2016- 17/MM/LTBS/CRPS/DOQ /S2/2915/2015	128/6-1	Kareepra	Kottarakkara	7.80 Ares	04/04/2016	03/04/2017	75000

2	Negarudeen, Charuvilaputhenveedu, Veliyam	2/2016- 17/MM/LTBS/CRPS/DOQ /S2/2877/2015	209/2	Odanavattom	Kottarakkara	5.13Ares	06/04/2016	05/04/2017	75000
3	Indirakumari, Palathodueedu, Neduvathoor	3/2016- 17/MM/LTBS/CRPS/DOQ /S2/1287/2015	254/6	Neduvathoor	Kottarakkara	9.50 Ares	11/04/2016	10/04/2017	75000
4	Surendran, Kalluvila, Karimpinpuzha, Pavithreswaram	4/2016- 17/MM/LTBS/CRPS/DOQ /S2/54/2016	65/3	Pavithreswaram	Kottarakkara	8.10Ares	11/04/2016	10/04/2017	75000
5	Pushppakumar, Uthradam, Nellimukku	5/2016- 17/MM/LTBS/CRPS/DOQ /S2/476/2016	295/11/3	Pallimon	Kollam	6.07 Ares	28/04/2016	27/04/2017	75000
6	Shaji, Anandam, Poothakulam	6/2016- 17/MM/LTBS/CRPS/DOQ /S2/380/2016	339/20-6	Poothakulam	Kollam	5.30 Ares	02/05/2016	01/05/2017	75000



7	Karim, Kunnelputhenveedu, Perumbuzha	7/2016- 17/MM/LTBS/CRPS/DOQ /S2/536/2016	46/8	Melila	Kottarakkara	9.61 Ares	04/05/2016	03/05/2017	75000
8	Manoshkumar, Ambikavilasam, Puthenkulam, Karimbaloor	8/2016- 17/MM/LTBS/CRPS/DOQ /S2/2024/2015	41/11	Parippally	Kollam	5.00 Ares	10/05/2016	9/05/2017	75000
9	L.Vijayan, Rajeshbhavan, Karuvelli.P.O	9/2016- 17/MM/LTBS/CRPS/DOQ /S2/612/2016	23/1-8	Valakom	Kottarakkara	9.98 Ares	13/5/2016	12/5/2016	75000
10	Sabumon, Kozhikkunnathuveedu, Kottathala	10/2016- 17/MM/LTBS/CRPS/DOQ /S2/1138/2016	325/72	Neduvathoor	Kottarakkara	4.10Ares	27/6/2016	27/6/2017	75000
11	Negarudeen, Charuvilaputhenveedu, Veliyam.P.O	11/2016- 17/MM/LTBS/CRPS/DOQ /S2/1356/2016	2/2-1	Ummannoor	Kottarakkara	4.10 Ares	18/7/2016	17/7/2017	75000

12	Sudarsanan, Mukaluvilaveedu, Punakkannoor. Alummoodu.P.O, Mampuzha	12/2016- 17/MM/LTBS/CRPS/DOQ /S2/1018/2016	86/4-3, 86/4	Odanavattom	Kottarakkara	9.72Ares	18/7/2016	17/7/2017	75000
13	RajendraPrsad, Chikkubhavan, Pallimon.P.O, Kollam	13/2016- 17/MM/LTBS/CRPS/DOQ /S2/540/2016	345/1	Veliyam	Kottarakkara	9.50 Ares	21/7/2016	20/7/2017	75000
14	Sivankutty Pillai, Cheppallil, Patharam.P.O	14/2016- 17/MM/LTBS/CRPS/DOQ /S2/2918/2016	424/22	Sooranadu South	Kunnathoor	Ares9.57	8/8/2016	7/8/2017	75000
15	Dinesh, Arappura Veedu, Vendar.P.O, Puthoor	15/2016- 17/MM/LTBS/CRPS/DOQ /S2/3013/2016	260/17	Kulakkada	Kottarakkarra	9.30Ares	8/8/2016	07/8/2017	75000
16	Karunakaran Unnithan, Vrundavanil, Madandankode	16/2016- 17/MM/LTBS/CRPS/DOQ /S2/162/2016	662/2	Kareepra	Kottarakkara	7.18 Ares	8/8/2016	07/8/2017	75000

17	Thulaseedharan, Chithirayil, Neeleswaram	17/2016- 17/MM/LTBS/CRPS/DOQ /S2/2036/2016	482/9-2	Naduvathoor	Kottarakkara	20.00 Ares	18/8/2016	17/8/2017	150000
18	Negarudeen, Charuvilaputhenvee du, Veliyam West.P.O	18/2016- 17/MM/LTBS/CRPS/DOQ /S2/1293/2016	35/9	Odanavattom	Kottarakkara	8.09 Ares	24/9/2016	23/9/2017	75000
19	L.Vijayan, Rajeshbhavan, Karuvilil.P.O, Ezhukone	19/2016- 17/MM/LTBS/CRPS/DOQ /S2/613/2016	23/6	Valakom	Kottarakkara	9.00 Ares	7/10/2016	06/10/2017	75000
20	Ani.S, Thekkadeth Veedu, Mukkoodu.P.O, Karippuram	20/2016- 17/MM/LTBS/CRPS/DOQ /S2/1243/2016	249/4-3	Pavithreswara m	Kottarakkara	9.20 Ares	17/10/2016	16/10/2017	75000

<b>Sl. No</b>	<b>Concession holder's name and address</b>	<b>Concession no.</b>	<b>Mineral</b>	<b>Survey no</b>	<b>Village</b>	<b>Taluk</b>	<b>Jaw size</b>	<b>valid from</b>	<b>valid to</b>
1	Shri.M.K.Biju, Karthika, Kuttikkad.P.O, Kadackal	44/2016-17/RMCU/KLM/2694/M3/2016,	Granite Building Stone	313/6, 6-2,6-3,6-4,312/7-1 & 7-2	Ittiva	Kottarakkara	(48"x9") 1200MMx225MM -2 Primary Crusher 30"x15" (750MMx379MM) - 1	28/3/2016	31/3/2017
2	Shri.M.K.Biju, Karthika, Kuttikkad.P.O, Kadackal	43/2016-17/RMCU/KLM/2695/M3/2016	Granite Building Stone	158/8, 158/6, 158/7-2, 158/7-1, 158/12-1, 158/12-2	Ittiva	Kottarakkara	Primary crusher 43"x27" - 1, VSI - 1	28/3/2016	31/3/2017
3	Shri.M.Abraham, Charuvila, Puthenveedu, Ayoor	151/2016-17/RMCU/KLM/3472/M3/2016,	Granite Building Stone	153/2	Valakom	Kottarakkara	1.40cmx22.50cm(16"x9")-1, Primary 24"x14"-1	04-08-2016	03-06-2017
4	M/s.Poabs Enterprises Ltd, Kuttoor.P.O, Thiruvalla	30/2016-17/RMCU/KLM/2567/M3/2016	Granite Building Stone	297/3, 297/4, 297/2	Chithara	Kottarakkara	cone crusher (215HP)-2 Primary (c-125), VSI - 1	23/3/2016	31/3/2017

## References

1. District Resource Map, Kollam district, Kerala. Published by Geological Survey of India.
2. Detailed Information on Bauxite in India Geological Survey of India. 1994. Retrieved from [http://www.portal.gsi.gov.in/gsiDoc/pub/DID\\_Bauxite\\_WM.pdf](http://www.portal.gsi.gov.in/gsiDoc/pub/DID_Bauxite_WM.pdf)
3. Rajan, T. N. and Anil Kumar, P. S. 2005. Geology and Mineral Resources of the States of India Part IX – Kerala. Geological Survey of India Miscellaneous Publication No. 30.
4. Mini Chandran 2013. Ground Water Information Booklet of Kollam district, Kerala. Technical Reports: Series ‘D’. Central Ground Water Board.
5. Retrieved from [kerenvis.nic.in](http://kerenvis.nic.in)
6. Retrieved from [keralasoils.gov.in](http://keralasoils.gov.in)
7. Retrieved from [dmg.kerala.gov.in](http://dmg.kerala.gov.in)

## **Geology of Kerala**

### **Physiography**

Physiographically the state can be divided into four domains from east to west, viz., the Western Ghats, the foothills, the midland and the coastal low- land.

### **Western Ghats**

The hill ranges of the Western Ghats rise to an altitude of over 2500m above the MSL and the crest of the ranges marks the inter-state boundary in most of the places. A breach in the continuity of the ranges marks the Palghat Gap with a sinistral shift of 50 km between the shifted crests. The Wynad plateau and the Munnar (10°57'00": 77°31'00") upland fall within this zone.

### **Foothills**

The foothills of the Western Ghats comprise the rocky area from 200 to 600m.above MSL. It is a transitional zone between the high -ranges and midland.

### **Midland region**

This forms an area of gently undulating topography with hillocks and mounds. Laterite capping is commonly noticeable on the top of these hillocks. The low, flat-topped hillocks forming the laterite plateau range in altitude from 30-200m and are observed between coastal low-land and the foothills.

### **Coastal low- land**

Coastal low-land is identified with alluvial plains, sandy stretches, abraded platforms, beach ridges, raised beaches, lagoons and estuaries. The low- land and the plains are generally less than 10m above MSL.

### **Rivers**

Kerala is drained by 44 rivers, many of which originate from the Western Ghats. Except Kabini, Bhavani and Pambar which are east - flowing, the rest of rivers are west- flowing and join the Arabian Sea. A few of them drain into the backwaters. Most important rivers (with their length in km in paranthesis) of the state, are Chandragiri(105), Valapatnam (110), Achankovil (120) Kallada (121), Muvattupuzha (121), Chalakudy (130), Kadalundi(130), Chaliyar (169), Pampa (176),Bharathapuzha (209) and Periyar (244).

### **Geology**

Geologically, Kerala is occupied by Precambrian crystallines, acid to ultra basic intrusives of Archaean to Proterozoic age, Tertiary (Mio-Pliocene) sedimentary rocks and Quaternary sediments of fluvial and marine origin (Fig.I). Both the crystallines and the Tertiary sediments have been extensively lateritised.

Based on the detailed studies by GSI during the last three decades, the following stratigraphic sequence has been suggested.



	Quaternary (Q)	Pebble bed Kadappuram Formation (marine) Periyar Formation (fluvial) Viyam Formation (fluvio-marine) Guruvayur Formation (Palaeo-marine) Laterite
	Mio-Pliocene (N 12) (Tertiary Tt)	Warkalli Formation (Sandstone and clay with lignite intercalations) Quilon Formation (Fossiliferous limestone and calcareous marl).
	Mesozoic (61-144Ma.)	Gabbro / Dolerite dykes
P	Younger granites (550-390Ma)	Alkali granites, granite, Granophyres and other acid intrusives
R		
O	Charnockites (younger) (550Ma)	Massive charnockite, incipient charnockite, Cordierite charnockite
T		
E	Ultrabasic/basics (Younger) (700-600Ma)	Perinthatta anorthosite, Kartikulam gabbro, Adakkathodu gabbro, Begur diorite
R		
O	Basic Intrusives (2100-1600Ma)	Agali- Anakkatti dykes
Z		
O	Migmatite/gneiss/older granitoid (PGC II) (Ptm) (2500-2200Ma)	Gamet-biotite - gneiss with associated migmatites, quartz-felspathic gneiss, homblende gneiss, homblende-biotite gneiss, quartz-mica gneiss
I	Vengad (A Dtv) Group	Quartz-mica schist and quartzite, conglomerate
A	Charnockite (older) (Ac) 2600Ma	Mafic granulite, pyroxene granulite, Banded magnetite quartzite and gneissic charnockite
R		
C	Khondalite Group (Ak)	Quartzite, mafic granulite, calc-granulite gamet- biotite-sillimanite-cordierite gneiss, gamet-biotite- gneiss, leptynite
H	Peninsular Gneissic Complex (PGC I) (Ap) (3000Ma)	Foliated granite, homblende gneiss, pink granite gneiss, biotite gneiss
A	Layered ultrabasic - basic Complex (3100- 3000Ma)	Peridotite, dunite, pyroxenite, anorthosite
E		
A	Wynad Schist Complex (Aw) (3200Ma)	Talc-tremolite schist, fuchsite quartzite, amphibolite, calc granulite, quartz sericite schist, kyanite quartzite, gamet - sillimanite gneiss/ schist, magnetite quartzite, kyanite mica schist
N		

Base not recognised

## **The Archaeans**

Rocks of Archaean Era encompass a wide spectrum of litho-assemblages ranging from khondalite, charnockite, gneiss and meta-sedimentary rocks occupying the Western Ghats including the foothill region. The Khondalite and Charnockite Group are correlated with the Eastern Ghat Supergroup based on the overall similarity in lithology and geochronology.

## **Wynad Supracrustals**

The meta-sedimentary, and ultramafic rocks occurring in the Wynad District generated keen interest among the GSI geologists in 1970s. The high-grade Wynad supracrustal rocks are correlated with the Sargur Schist Complex of the Karnataka (Nair, *et al*, 1975; Adiga, 1980). The schistose rocks are characterised by intense deformation, medium to high-grade metamorphism, migmatitisation and lack of sedimentary structures. The schist complex consists of meta-ultramafites, schist, meta-pelites, meta-pyroxenite, serpentinite, talc-tremolite rock and amphibolite.

The meta-sedimentaries occur as thin linear bodies within the migmatites. These consist of pelites, psammopelites and quartzites. The predominant rock types are corundum- mica schist, kyanite schist, quartz- mica schist and iron stone.(Anil Kumar *et al*,1993).

These rocks occur as narrow arcuate belts, lenses, and other forms of enclaves within Peninsular gneisses and charnockite. The group can be divided into medium-to low- grade metasedimentary rocks and meta-mafic and meta-ultramafic rocks. The lithology of the high-grade schists consist of quartz-mica schist with kyanite, quartz-sericite schists,quartzites,magnetite quartzite, fuchsite quartzite and meta-ultramafites. Their contact with the surrounding gneisses are concordant due to later co-folding. Several linear belts of such high-grade schists and ultramafite enclaves occur as isolated bands within the granulite and gneissic terrain of Kasaragod and Kannur districts.

## **Layered ultrabasic- basic complex**

Remnants of layered basic- ultrabasic complex are reported from Attappadi area(Nambiar 1982).The ultramafics are represented by meta-pyroxenite, meta-dolerite, peridotite with chromite and meta-gabbro (Lahiri *et al*, 1975). The anorthosite of Attappadi is only a few metre thick. Occurrences are around Narsimukku, Pudur and Melmulli areas. An east-west trending narrow lenticular body of serpentinitised dunite is reported from Punalur mica mine belt.

A minor body (200 m long and 10-15m wide) of anorthosite was reported within pyroxene-granulite-charnockite terrain from Katanjari *parambu* of Kasargod district (Adiga, 1979).

Another dismembered layered igneous complex consisting of alternate layers of peridotite and pyroxenite within charnockite was traced around Panathadi area of Kannur District (Adiga, 1980).

### **Peninsular Gneissic Complex- I (PGC-I)**

The rocks of Peninsular Gneissic Complex (PGC) are exposed in the northern parts of Kerala adjoining Karnataka. This consists of a heterogeneous mixture of granitoid materials. The equivalent rocks of PGC in Kerala include hornblende-biotite gneiss (sheared), biotite-hornblende gneiss, foliated granite and pink granite gneiss. Granite gneiss is exposed along the intra-State boundary of Palakkad District as well as in Idukki District. Pink granite gneiss, though widespread, is best developed in Devikolam (10°04'00": 77°06'30"), and Udumbanchola (10°00'00":77°15'00") areas of Idukki District.

This consists of gneisses showing preponderance of either hornblende or biotite. The percentage of hornblende and biotite varies from place to place. This can be traced from Manantoddy to further northwest upto the west coast. West of Manantoddy, the rock is hornblende gneiss. It shows coarse granulitic to gneissic texture and is composed of hornblende, feldspar, quartz, pyroxene, biotite and garnet. Alkali feldspar shows alteration to clay and sericite. Biotite is mainly secondary after hornblende.

Around Mahe and Thalasseri, the biotite gneiss (Nair *et al.*, 1974) is medium-grained and gneissose rock consisting of alternate layers of mafics and felsics.

### **Khondalite Group**

The Khondalite Group of rocks include calc-granulites, quartzite and para-gneisses of pelitic parentage. Para-gneisses are ubiquitous and are well-developed in the southern part of the state, particularly, in Thiruvananthapuram and Kollam districts. Calc-granulite and quartzite occur as bands within the para-gneisses and amidst the Charnockite Group and migmatitic gneisses.

#### ***Calc-granulite***

Calc-granulite occurs as linear bands mainly in the eastern part of Kollam and Thiruvananthapuram District, northeast and east of Munnar in Idukki district and in parts of Palakkad District. The rock is generally medium to coarse-grained, inequigranular and granoblastic in texture. It consists of diopside and plagioclase. Minerals like wollastonite, scapolite, calcite, garnet, spinel, sphene, quartz and apatite are also present in different proportions.

### ***Quartzite***

Quartzite occurs as linear bands amidst the khondalitic gneiss, charnockite and migmatitic gneisses. These bands are exposed between Pathanamthitta (9°15'45": 76°47'00"), and Muvattupuzha (9°59'00": 76°35'00") in Ernakulam District. The rock is coarse-grained and generally white in color with a brownish coating on the weathered surface. It consists of granular quartz with subordinate feldspar, garnet and iron oxide.

### ***Garnetiferous biotite-sillimanite gneiss***

Garnetiferous biotite- sillimanite gneiss is well-developed in the southern part of the state. It occurs in close association with the migmatitic gneisses, charnockite and charnockite gneisses, mostly as weathered outcrops. Sillimanite- rich bands occur alternating with garnet - rich portions or with quartzo-feldspathic layers. Rutile and iron oxides are the common accessory minerals.

### **Charnockite Group**

Charnockite Group shows great diversity in lithology comprising pyroxene granulite, hornblende pyroxenite, magnetite quartzite, charnockite and hypersthene-diopside gneisses and cordierite gneiss. Charnockite and charnockitic gneiss have preponderance over all other crystalline rocks covering 40 -50% of the total area of the State. The charnockites are well-exposed in the central and northern parts of Kerala including the high-hills of the Western Ghats. Charnockite has lesser predominance in Thiruvananthapuram and Kollam districts. In Attappady, the Bhavani Shear Zone is limited by the charnockite massif of the Nilgiri plateau on the north. Though the interrelationship of the Charnockite and the Khondalite is not clear, in many places there are intercalations rather than interlayering of one with the other. In Palakkad District, the Khondalite Group of rocks structurally overlies the charnockite. The occurrence of pyroxene granulite as fine and linear bodies within the charnockite of Tirur, suggests that charnockite is a product of migmatization of pyroxene granulite (Vidyadharan and Sukumaran, 1978). Charnockite and charnockitic gneiss consist of quartz, feldspar and biotite. Garnet-bearing variants are also observed. The basic charnockite is more granulitic and contains clino- and ortho- pyroxenes, feldspar, biotite and garnet whereas the acid variety (alaskite/ enderbite) is greenish black, coarse-grained, massive to poorly foliated rock consisting of quartz, feldspar and pyroxenes. Basic charnockite has low- potash feldspar and more clinopyroxene. This is devoid of garnet and graphite, but shows a little amount of biotite (Chacko, 1922). Due to the polygenetic nature of the rock, geochemical and

mineralogical variations do exist between charnockites reported from Kerala. In the Periyar valley region, in Idukki and Kottayam districts, pyroxenite and alaskite constitute the Charnockite Group (Nair, and Selvan, 1976).

The available age data indicate that the massive charnockites are older and their ages range between 2155 and  $2930 \pm 50$  Ma (Soman, 1997).

Also charnockite has been subjected to retrogression and migmatization.

## **Archaean to Palaeo-Proterozoic**

### **Vengad Group**

A succession of schistose rocks in parts of Tellicherry taluk in Kannur district is described as Vengad Group of rocks (Nair, 1976). The Vengad Group comprises of basal conglomerate, quartzite and quartz-mica schist. The contacts are highly gradational. The conglomerate shows graded bedding and quartzite shows current-bedding.

An angular unconformity marked by conglomerate horizon extending from Kuthuparamba ( $11^{\circ}49'30''$ :  $75^{\circ}34'00''$ ) to Vengad ( $11^{\circ}53'30''$ :  $75^{\circ}32'00''$ ) in Kannur district, separates the younger quartz-mica schist and quartzite from the older schistose and gneissic rocks. The lithology consists of basal oligomictic conglomerate, quartzite, quartz-biotite-muscovite schist and biotite quartzite. The schists are exposed over an area of 300 sq km having a lensoidal shape with its longer axis trending in NW-SE direction. The basement rock is gneissic or migmatitic with relicts of high-grade schists, ultramafites and quartzites of the Wynad Schist Complex. Four major occurrences of conglomerate are noticed in a NW-SE direction over a length of 10 km.

Lack of migmatization, presence of primary structures and low-grade metamorphic minerals characterize these rocks.

### **Migmatite\ Gneiss\ Granitoid (PGC-II)**

#### **Quartzo-feldspathic gneiss**

Migmatite includes variety of gneissic rocks which are next in importance to charnockite as a dominant litho-assemblage. Quartzo-feldspathic gneiss occurring along the contact zone between garnet-biotite gneiss and garnet-sillimanite gneiss of Thiruvananthapuram area represents an original intrusive phase. It is a feebly foliated, fine-grained, leucocratic granulitic

rock occurring in close association with garnet-sillimanite gneiss and garnet-biotite gneiss with gradational contact relationship in the southern parts of Kerala. The origin of this rock is attributed to stress-induced injection of acid materials into the host rocks (Nageswara Rao and Raju, 1970).

### **Garnet- biotite gneiss**

Garnet-biotite gneiss is well-developed in the northeastern parts of Kollam and Thiruvananthapuram districts. This carries inclusions of pyroxene granulite and disseminations of graphite at many places (Jacob, 1965). It consists of quartz, microperthite, biotite, plagioclase and graphite. This rock also occurs in the northern parts of Palakkad District in close association with khondalite, charnockite and hornblende gneiss. These rocks are subsequently formed by retrogression and migmatization of the Khondalite Group.

East of Kottayam and Idukki districts, light grey, pink garnet-bearing biotite gneiss is widely seen. It is a gneissic granulite. The presence of biotite and concentration of garnet in layers give the rock a banded appearance. (GSI, 1995).

### **Hornblende gneiss, hornblende-biotite gneiss, quartz- mica gneiss**

These rock types occur within the migmatites and associated retrograded charnockite. The naming is purely based on the preponderance of the minerals and these rocks occur in the Periyar valley area east of Thodupuzha. (Nair and Selvan, 1976). These medium-grained, foliated, banded rocks consist of alternate layers rich in hornblende or biotite. Bands of coarse to medium-grained light grey to pink granite traverse these rocks. Hornblende- biotite gneiss showing lit par lit relationship with the granite gneisses is the dominant rock type in the Periyar valley. This is admixed with contorted bands and enclaves of pyroxene granulite, calc-granulite and hornblende- biotite granulite. These are highly deformed.

In the Palakkad gap area, these gneisses occur over a large area, showing migmatitic structures such as agmatites, nebulites, schlierens, ptygmatic folds, quartzo-feldspathic neosomes and ferromagnesian palaeosomes. (Muraleedharan and Raman, 1989).

## **Proterozoic**

### **Basic intrusives**

Basic dyke emplacements within the Archaean crystalline rocks of Kerala are spread throughout the entire length and breadth of the state. Of these, dolerite dyke occurring north of the Palakkad gap had given Proterozoic age whereas in the south this dyke is of Phanerozoic

age. The older basic dykes are metamorphosed along with the country rocks and are now recognised as epidiorite and amphibolite. Another set of dykes, apparently post-dating the regional metamorphic event are subjected to thermal metamorphism. Clouding and sericitisation of feldspars and uralitisation of pyroxenes are common in such dykes. In the absence of chronological data such dykes are considered to be of Proterozoic age. Most of the dykes are vertical in disposition and are traced as linear features. *En-echelon* pattern of some dyke swarms suggests that magmatic intrusion was controlled by shearing of the host rock., Mineralogically, the dykes are made up mostly of plagioclase feldspar and pyroxene(augite and aegirine-augite) with magnetite, apatite and olivine as accessories. The ENE-WSW dolerite dyke swarm of Agali- Anakkati area in Palakkad District within the Bhavani Shear zone showed in isotopic age from 1900 to 2000 Ma (Radhakrishna and Mathew Joseph, 1993). The rock is highly jointed and altered (Jacob, 1965). Similar basic intrusive bodies are traced in the Achankovil shear zone in Vazhamuttam (9°14'00":76°46'40"), Kulasekhara pettah (9°16'00":76°47'45") (Thomas Mathai *et al*, 1984). Sheet-like bodies of fine to medium-grained, dark coloured meta-gabbro occurs in Periyamuli (11°13'00"; 76°43'00") for about 20 km in ENE-WSW direction, Karuvarai (11°04'00"; 76°32'30") and few gabbro bodies south of Thuvapattu (11°06'30"; 76°44'45") in Attapady valley, Palakkad district. Meta-gabbro forms small hillocks east of Payyanam (10°31'00"; 76°21'00"), southwest of Kainur (10°36'00"; 76°09'00") and Chemmannur (10°41'00"; 76°01'00"), Vaga (10°35'00"; 76°06'00") and Arthat (10°37'00"; 76°03'00") in Trichur District (Mahadevan, 1962).

Dykes in north Kerala show , NW-SE, NE-SW and NNW-SSE trends. Host rocks are charnockite, gneisses and supracrustals(Radhakrishna *et al* 1991). Dykes are mainly dolerite but occasional meta-gabbro or meta-norite are also traced. In Agali- Anaikatti area of the Attapadi- Bhavani shear zone, dykes are confined within a 20-25km wide zone and extend from west of Agali to eastward for about 100km following a ENE-WSW direction.(Radhakrishna, *et al*, 1999).

The rock consists of 95% calcic plagioclase, 5% clinopyroxenes and subordinate amounts of magnetite. There are a number of concordant and discordant basic intrusive of dolerite and gabbro,meta-gabbro, meta-norite, meta-pyroxenite and anorthositic gabbro. These are not mappable and are seen in Pappinpra (11°06'20", 76°05'56") Velli(11°04'00":76°07'45"),



Kalpetta (11°04'12":76°05'32"). An extensive basic diorite has been mapped over an area of 25 sq km at Panavalli (11°53'30",76 ° 2'30"; Nair, *et al* 1976).

The rock is composed predominantly of calcic plagioclase (95%) rest clinopyroxene with subordinate amount of magnetite. Another relatively small body of anorthosite is around Kalivalli (11°51'30"; 76°12'30") in south Wynad taluk, Wynad District.

### **Ultrabasic/ basic intrusive (younger)**

#### **Perinthatta Anorthosite**

A major elliptical body of anorthosite spread over an area of more than 50 sq.km is reported from Perinthatta (12°10'00":75°17'30";Vidyadharan *et al*, 1977). The anorthosite is with a very irregular border and a tongue-like projection into the country rock of charnockite and pyroxene granulite of Kannur District.The anorthosite is coarse to very coarse-grained, and shows variations from pure anorthosite to gabbroic anorthosite and gabbro from the centre to the periphery suggestive of zoning. The modal composition corresponds to nearly 95% plagioclase (An<sub>58-72</sub>) and <10% clinopyroxene, apatite, calcite and magnetite. The gabbroic variants have more of mafics.

The structural configuration suggests that the anorthosite was emplaced in synformal structure as a phacolith. The flow-banding in anorthosite indicates its syntectonic emplacement. The Perinthatta anorthosite is assigned a Proterozoic age.

#### **Ezhimala gabbro-granophyre complex**

The major high-relief feature proximal to the Perinthatta anorthosite is constituted by the gabbro-granophyre Complex (Nair and Vidyadharan, 1982). The granophyre massif is fringed by the gabbro to the east and south. The Bavali fault running north of the complex is presumed to have dismembered the body from the Perinthatta anorthosite. Locally, the gabbro has anorthositic differentiates within it. Veins of granophyre traverse the gabbro at places give rise to breccia-like structures. The granophyre shows a sharp contact with the gabbro into which it intrudes. Rapakivi structure is observed within the granophyre. According to Nair and Vidyadharan (1982) rocks of Ezhimala complex display bimodal character with conspicuous basic and silicic components.

#### **Kartikulam and Karraug Gabbro**

Two gabbro bodies namely Kartikulam gabbro and Karraug gabbro are located northeast of Manantoddy bordering the Karnataka (Nair *et al*, 1975).The gabbro body at Kartikulam

occupies an area of about 45 sq.km. with an elliptical shape within the gneissic terrain. The actual contact with the gneiss is concealed but it is believed to be sharp. At many places, the gabbro is agmatized by coarse quartzo-feldspathic material.

The gabbro is coarse-grained and of uneven texture consisting essentially of plagioclase and pyroxene. Variation to anorthositic composition is noticed. The plagioclase is of labrodorite composition and shows alteration to sericite at places (Rema Warriar and Venkataraman, 1986). The pyroxenes are uraltized to varying degrees.

The Karraug gabbro body is located east of it and south of the Kabini River. It shows similar features as that of the Kartikulam gabbro. The rock shows phenocrysts of feldspar set in a fine matrix of flaky minerals.

### **Adakkathodu gabbro**

At Adakkathodu (12°31'35"; 75°10'25"), northwest of Manantoddy, a 8 km long meta-gabbro, is intrusive into the basement gneisses on three sides and the Wynad schists in the east. It occurs proximal to the Bavali fault/lineament. It encloses, patches of quartz-sericite schists and biotite gneiss. (Nair *et al*, 1975). The rock is mesocratic to melanocratic, medium to coarse grained consisting mainly of pyroxene and plagioclase. The rock shows sub-ophitic texture and consists of enstatite and intermediate plagioclase of andesine-labrodorite composition (Nair *et al*, 1976). While the gabbro bodies of Kartikulam and Karraug to the east are olivine-bearing, the Adakkathodu gabbro is enstatite-bearing. Olivine, augite and zoned feldspars are recorded from the eastern body while the western body is enstatite-bearing, without the zoning in feldspar.

### **Begur diorite**

An extensive basic diorite body (25 sq.km.) has been traced north of Manantoddy in the Begur Reserve Forest (Nair, *et al*, 1976). It extends from Thirunelli to the Karnataka State border. The southern contact is with augen gneisses indicating emplacement along shear zones while the northern one with sillimanite gneisses. Aplite and dolerite veins are seen traversing the rock mostly parallel to the regional foliation. The rock is mesocratic to melanocratic, coarse-grained and consists of pink to grey feldspar, hornblende and biotite.

The rock is feebly gneissic and at places porphyritic (Rema Warriar and Venkataramana, 1986). The phenocrysts are mostly plagioclase. Mafics at times swerve round the phenocrysts

giving rise to augen structure. Hornblende is altered to biotite and chlorite. Accessories include epidote, apatite, zoisite and opaques.

The diorite shows tholeiitic characteristics. The diorite is considered as a transitional rock from the gabbro with which it is spatially associated in the nearby area with the plagioclase become more sodic.

### **Charnockites [younger]**

The area south of Palakkad exposes charnockite over large areas. The charnockites are represented by acid micropertitic charnockite and intermediate gneissic charnockite occurring in association with garnetiferous biotite gneiss and khondalite (Narayanaswamy and Purna Lakshmi, 1967). Massive charnockites are developed on a regional scale and occur as mappable litho-units (Raju and Gopalakrishnan, 1972), around Nedumangad. The massive charnockites in majority of the cases are acid and intermediate in composition. The rock is medium to coarse-grained and shows xenoblastic texture. It is composed of quartz, feldspar, pyroxenes, garnet and graphite with accessories like biotite, zircon, apatite and monazite.

Small patches, lenses or veins of charnockite occur in the gneisses of amphibolite facies in the Thiruvananthapuram area (Nageswara Rao and Raju, 1970). Here, the incipient charnockites are thought to have formed by transformation of paragneisses. (Hansen *et al*, 1987; Santosh *et al*, 1990). A few dominant varieties of incipient charnockites have been categorized by Ravindra Kumar and Chacko (1986) on the basis of their mode of occurrence, association and chemical processes involved in their development. At Kottavattom, north of Thiruvananthapuram, the charnockite consisting of quartz, K-feldspar, plagioclase, biotite, garnet and orthopyroxene as essential minerals and graphite, zircon, ilmenite, monazite, apatite, rutile and magnetite as accessory minerals are products of transformation of gneisses into coarse-grained charnockites along a system of conjugate fractures and foliation planes. (Saritha and Santosh, 1996).

### **Cordierite or Charnockite Gneiss**

Cordierite bearing large linear zones of charnockites were reported around Pathanamthitta (Nageswara Rao and Jacob, 1967) area. Cordierite charnockites or orthopyroxene-garnet-cordierite bearing gneisses (Sinha Roy *et al*, 1984; Santosh, 1987) occur as discontinuous bodies in the northern parts of Thiruvananthapuram and in selected stretches further south around Koliakode. The rock is composed of cordierite, orthopyroxene, plagioclase, K-feldspar, spinel and quartz and a little garnet and biotite.

The growth of cordierite and orthopyroxene took place concomitantly during the conversion of gneisses to charnockites. At Nellikala in Pathanamthitta, the cordierite occurs as anhedral grains of variable sizes in the charnockites (Nandakumar, 1996).

### **Younger granites**

The granites and its variants occur around Chengannur in Alappuzha and Pathanamthitta districts, Munnar in Idukki District, Peralimala in Kannur district and Kalpetta and Ambalavayal in Wynad District. Many of these granites occur as later emplacements along crustal fractures and faults. The Achenkovil – Tamraparni tectonic zone, the Attapadi shear zone, Bavali shear zone and the Moyar shear zone are all marked by granitic emplacements

### **Ambalavayal granite**

The Ambalavayal (11°37'15"; 76 °03'30") granite having an oval shape covers an area of 50 sq.km. The granite is light pink in color and is composed of quartz, pink feldspar, hornblende and biotite. The pegmatites traversing the granite show occasional flakes of molybdenite. The Amabalavayal granite occurring in the proximity of the Bavali lineament is thought to be emplaced during its reactivation. The granite is intrusive into the hornblende-biotite gneiss (migmatite) and the Wynad Supracrustals (Anilkumar *et al*, 1993). Four types of granites are recorded, viz., foliated granite, pink granite, grey granite and aplitic granite.

The foliated granite consists of quartz, microcline, orthoclase, plagioclase, biotite, hornblende, chlorite, calcite and zircon. The pink granite is a medium-grained consisting of quartz, microcline, plagioclase, sericite, chlorite, apatite, rutile, zircon and biotite. The grey granite is a medium-to fine-grained rock consisting of quartz, microcline, sericite, biotite, chlorite and calcite. The aplitic granite is a very fine-grained massive rock consisting of quartz, microcline, orthoclase, plagioclase, sericite, biotite, calcite, chlorite, apatite and opaques.

K-Ar age of Ambalavayal granite ( $560 \pm 30$  Ma, Nair, *et al*, 1985) is lower than Rb-Sr age ( $595 \pm 20$  m.a Santhosh *et al*, 1986), but is higher than that of U-Pb-age ( $505 \pm 20$  ma, Odom, 1982). The reason for this variation in the date may be attributed to the different techniques adopted and also to the presence of biotite of multiple generation.

### **Munnar granite**

The Munnar (10°05'00"; 77°05'00") granite with an areal extent of 50 sq km is an E-W trending irregular body emplaced within the migmatite and apophyses extend into the surrounding gneisses. The granite dated to be  $740 \pm 30$  m.y (Odom, 1982) is traversed by pegmatite, aplite

and quartz veins. Three types of granite are recorded. Foliated granite, Coarse pink granite and medium grey granite. The foliated granite consists of stringers and streaks of mafics consisting of biotite, hornblende, chlorite and magnetite alternating with felsics consisting of quartz and potash feldspar. Potash feldspar is predominantly orthoclase. The closely spaced foliations are persistent but discontinuous. This granite forms a domal structure south of Munnar. It has a sharp contact with the migmatite. Coarse pink granite consists of pink feldspar, quartz and a little amount of mafics. Mafics are biotite, sphene and hornblende. Medium grained grey granite, consists of quartz, feldspar, biotite, chlorite, zircon, sphene, epidote, calcite and sericite.

Major element data of Munnar granites do not show any significant variation amongst the three granites. Content of iron is more in medium grey granite and foliated granite. Different variation diagrams reveal a slight tendency towards alkali granite. The foliated granite shows more percentage of orthoclase than the other two granites. (Nair and Anil Kumar, 1990)

### **Ezhimala granophyre – granite complex**

A prominent granophyre body forms the hill known as Ezhimala, covering an area of 20 sq km in Kannur District. The granophyre is associated with gabbro and granite and is traversed by dolerite dykes. Two types of granophyres have been deciphered; coarse-grained leucocratic one and medium-grained one with more mafics. Drusy type, confined to higher elevation contain numerous vug lines with secondary minerals like quartz and calcite. Rocks of Ezhimala Complex display bimodal character with conspicuous basic and silicic components and total lack of rocks of intermediate composition typical of anorogenic suites (Nair and Vidyadharan, 1982). The granophyre is pink to ash grey coloured, massive, fine to coarse-grained, holocrystalline with equigranular texture. The granites are of two types. The major light pink granite with less of mafics show gradational relationships with the more greyish porphyritic variant (Varadan and Venkataraman, 1976).

Granophyre shows a typical granophyric intergrowth of quartz and feldspar forming the ground mass with phenocrysts of potash feldspar and some zoned plagioclase. The groundmass is totally of orthoclase. Augite is the chief ferro-magnesium mineral. Accessories include apatite, sphene, epidote, calcite and magnetite. Texturally the rock shows variation from coarse-grained leucocratic types with less mafics in the southern portion of the hill and medium to coarse grained type towards northern parts.

Minor outcrops of rapakivi granites are recorded within the granophyres of Ezhimala Complex. Anorthosites of Perinthatta and Kadannappally and granite, granophyre of Ezhimala together form the Ezhimala Complex. The light pink granite with less mafics is the major variety showing a gradational relationship with the more greyish porphyritic variety. The porphyritic variety, at places, shows rapakivi structure. The porphyritic granite shows mantled feldspar megacrysts. This variety grades into porphyritic granites without mantled feldspar and at higher levels grades into granophyre. The granite contains 60% of orthoclase feldspar, 5-10% of plagioclase, 20-25% of quartz with 4% of biotite, epidote, magnetite and fluorite. The low initial  $Sr_{87}/Sr_{86}$  ratio indicate that the rocks have a relatively minor amount of older sialic material. The Rb-Sr age of the granophyre is estimated to be 678 m.y (Nair and Vidyadharan, 1982). The Ezhimala Complex lies in close proximity to the Bavali lineament suggesting reactivation along the lineament and intrusion of the body.

### **Kalpatta granite**

The Kalpatta ((11°36'15";76°05'15")) granite is an oval- shaped intrusive into the Wynad schist and covers an area of 44 sq km (Rao and Varadan, 1967). The rock is grey coloured, medium-grained, homogenous biotite granite and has sharp contact with the country rock. A feeble foliation is imparted to the granite at places by biotite flakes. Xenoliths of amphibolite / hornblende gneiss are visible near the periphery. Irregular veins of pegmatite / aplite traverse the granite and also the enclaves. The K-Ar age of the biotite from the Kalpatta granite is dated as  $512 \pm 30$  m.a (Nair *et al*, 1985) and 527 m.a (GSI). Presence of enclaves and absence of significant replacement textures along with the geochemical characteristics assign a magmatic parentage for the granite. The proximity of the pluton to the Bavali lineament probably suggests intrusion along this fracture.

Three types of granites such as coarse grained biotite-granite, fine grained biotite granite, and porphyritic granite are mapped on the basis of texture, colour and mode of occurrence. Coarse-grained granite is a massive bluish grey rock with large xenoblasts of quartz and feldspars. The accessories include biotite, zircon, apatite and sphene. Blastesis of feldspar and sphene are common. Microcline, orthoclase, and plagioclase are seen as the major feldspar. Plagioclase composition varies from albite to oligoclase. This rock is exposed in Trikkaiappetta (11°35'04":76°08'41":), Manikkunnu (11°35'41":76° 07'09"), Kuttamangalam (11°30'08":76°07'11":) (Anilkumar *et al*, 1993).

Fine biotite-granite is a fine grained massive rock exposed around Muttimala (76°06'38":11°37'06"). It consists of orthoclase, quartz, microcline, biotite, sericite, zircon, sphene, apatite and opaques. Myrmekitic quartz is recorded. Pophyritic granite consists of myrmekitic quartz, microcline, sericite and biotite. Very coarse grained biotite with included crystals of orthoclase, microcline and albite are common. Except for the texture, all the three granites show similar characters. ( Anilkumar, *et al* 1993). Based on Rb-Sr dating , Kalpatta granite is dated 765 Ma. (Odom 1982).

### ***Chengannur granite***

The Chengannur (9°18'45"; 76°31'00") granite in Pathanamthitta District is an oval shaped body with the long axis trending in east-west direction covering an area of 15 sq.km in and around Chengannur. The granite is intrusive into the charnockite gneisses. The body is emplaced close to the Achankovil shear zone. K-Ar date of the hornblende indicates an age of 550 m.a (Soman *et al*, 1983). The Chengannur granite is inferred to be a post kinematic granite of magmatic parentage.

Two types of granites are recorded. One is medium-grained pink granite and the other is coarse-grained grey granite. The former consists of quartz, perthitic feldspar, plagioclase, biotite, hornblende, apatite and zircon. The composition of plagioclase varies from albite to oligoclase. Microcline perthite is also seen. The coarse grained grey granite consists of perthite, plagioclase, hornblende, biotite, quartz with occasional occurrence of hypersthene,apatite and zircon. Hornblende and biotite are less common by occurred minerals than hypersthene. Relicts of hypersthene are also seen. This granite may be a product of granitisation of charnockite. K<sub>2</sub>O content always exceeds that of Na<sub>2</sub>O . The high SiO<sub>2</sub>,high alkali, high Fe/Mg ratio, high values of Gallium indicate that the granite belongs to alkali type. It might have an origin from recycled and rehydrated continental crust. ( Nair and Anil Kumar,1990).

### ***Peralimala granite***

The Peralimala (11°09'19":75°38'46") alkali granite is a linear intrusive body emplaced along the axial trace of a mega fold in EW direction. Peralimala intrusive body occurs as a diatreme of alkali composition with a maximum linear extension of 15 km and a width of 3 km. Based on colour, texture, composition and mode of occurrence four types of granites are identified. These are pink gneissic granite, porphyritic granite, grey granite and pink granite. Pink alkali granite is a coarse-grained rock consisting of microcline, orthoclase, plagioclase, quartz,



hornblende, epidote, aegirine, sphene, calcite, perthite and apatite. Quartz is present in only subordinate amounts. Feldspar content is very high. The preferred orientation of feldspar gives a crude alignment. At Perumpunna, (75°44'00":11°55'28") pink gneissic granite shows preferred orientation of biotite and pyroxene. The porphyritic granite occurs as a lensoidal body containing quartz, feldspar, pyroxene and hornblende. Feldspar forms the phenocrysts in a matrix of quartz-feldspar and mafics. Grey granite is a coarse- to medium- grained rock with microcline, quartz, orthoclase, perthite, hornblende and zoisite. Light grey granite is a medium-grained rock consisting of microcline, orthoclase, plagioclase (albite to oligoclase), epidote, aegirine, hornblende and rutile. The major element chemistry of the granite do not show much variation. The pink granite shows high content of potash. A negative correlation for K<sub>2</sub>O content with respect to SiO<sub>2</sub> is very pronounced for pink granite owing to its alkaline nature. Barium and strontium show very high values for Peralimala granite. ( Anilkumar *et al*,1993).

### **Sholayur granite**

The Sholayur (11°04'15";76°42'00") granite, is exposed around Kuttiyadikal Mala (11°01'52":76°42'00") and Vachchpathi (11°04'15":76°44'00"). It is a homophanous medium-grained, pink coloured granite, consisting of quartz, orthoclase, microcline, oligoclase, perthite, aegirine augite, biotite, hornblende and sphene. In some places, calcite, apatite, sericite are also observed. The schlierens mark the contact zone of the granites with the host rock. This granite is emplaced within the Wynad supracrustals. SiO<sub>2</sub> varies from 58.76 to 73%, Al<sub>2</sub>O<sub>3</sub> 14% to 17%, Na<sub>2</sub>O 1.8% to 2.4% and K<sub>2</sub>O 0.8 to 1.5%. The distribution of SiO<sub>2</sub> is highly non-uniform within the same type of granite. The pink granite is becoming alkali granite at places.(Anil Kumar and Nair,1992).

### **Intermediate intrusives**

The syenite body at Mannapra (10°30'00";76°32'00") is exposed as an elongated NW-SE trending body covering an area of 8 sq km in Thrissur District. The syenite intrusive, makes sharp contact with the charnockite near the charnockite-migmatite contact. The rock is medium to coarse- grained at its peripheries and tends to be coarse-grained towards the centre. Mineralogically, the rock is composed of alkali feldspar, orthopyroxene, clinopyroxene and amphibole with minor amounts of plagioclase, biotite and opaques. A small syenite (Angadimugar syenite) body is located in Kumbala village (12°35'15"; 76°07'00") and about 20 km east of Kumbala in Kasaragod District. The intrusive body has an elliptical outline and covers an area of 5 sq km. The body is intrusive into the Khondalite Group and encloses



enclaves of amphibolite in the peripheral parts. The rock is medium to coarse grained, light grey and massive.

### **Mesozoic intrusives**

#### **Basic intrusives**

Basic intrusives in Kerala, mainly represented by dyke swarms in NNW-SSE to NW-SE trend, cut across all the metamorphic rocks and the earlier structural trends. Their unmetamorphosed nature and stratigraphic relation with the country rocks prompted their correlation to the Deccan Trap volcanism.

The basic dykes have been emplaced into the migmatites and charnockite in NNW-SSE to NW-SE and ENE-WSW directions along distensional and shear fractures respectively. Dolerite dykes of Kerala are mostly quartz tholeiites rarely clinotholeiite. The basic dykes of Pathanamthitta ( $9^{\circ}15'45''$ : $76^{\circ}45'30''$ ) are genetically unrelated types. These dykes have not undergone any internal differentiation during intrusion.

The variation in the chemistry of individual dykes may be due to the cogenetic differential sequence. Dolerite dykes intrude the country rocks at an angle greater than  $80^{\circ}$ . The dolerite dykes of Kuttuparamba ( $11^{\circ}49'30''$ : $75^{\circ}34'00''$ ) in Kannur District shows cross cutting relationship with all the formations. The basic dykes of Vamanapuram ( $8^{\circ}43'00''$ : $76^{\circ}54'00''$ ) are either gabbroic or doleritic intruding the gneissic rocks. These are trending NNE to SSW and NNW to SSE directions and are unmetamorphosed. Mineralogically all these dykes show more or less same composition except the meta-dolerites. Variation in the trace elements like Ti, Zr can be attributed to the differential degree of partial melting of the mantle material. (Nair and Gopala Rao, 1989).

The unmetamorphosed Idamalayar gabbroic dyke with a NNW-SSE trend is traced for over 80 km in the central part of Kerala. The rock is mesocratic, medium-grained, porphyritic and is composed of plagioclase (andesine to labradorite), hornblende and opaques. The reported age of 75 m.y for the Idamalayar dyke (Subramaniam, 1976) links it in time-relationship with Deccan Trap volcanism.

The NNW-SSE trending leucogabbro dykes in central Kerala dated by whole rock K-Ar method gave an age of  $81 \pm 2$  m.y and the NW-SE trending dolerite dyke  $69 \pm 1$  m.y. The

dolerite dykes are thought to have represented the feeder system for Deccan Trap volcanic sequences (Radhakrishna *et al*, 1994).

Basic dykes of Pathanamthitta area yielded ages of 99 Ma to 117 Ma and there are dykes which have yielded ages  $104 \pm 5$  Ma,  $127 \pm$  Ma and  $476 \pm$  Ma. These wide variations may be due to a protracted history of emplacement and the effect of Eocambrian to palaeozoic tectonothermal events affecting this region (Sinha Roy and Ramakrishnan, 1983.)

In Thiruvananthapuram District, Anakudi and Nedumannur dolerite dykes are dated by K-Ar method and the whole rock ages are  $104 \pm 5$  Ma and  $127 \pm 2$  Ma respectively (Sinha Roy and Ramakrishnan, 1983).

### **Tertiary Sedimentary rocks**

Mio-Pliocene sedimentary rocks are fairly widespread in the southern coastal belt, their remnants being noticeable in the central and northern coastal areas. These sedimentary rocks consist of a series of variegated clay and sandstones with lenticular seams of lignite, known as Warkalli Formation, underlain by more compact marly sands with shell fragments and thin horizons of limestone (Quilon Formation).

The Tertiary sediments have a gentle dip towards west. The Warkalli Formation extends in a narrow belt from Thiruvananthapuram ( $8^{\circ}28'30''$ :  $76^{\circ}57'20''$ ) to Kasaragod ( $12^{\circ}30'00''$ :  $74^{\circ}59'00''$ ) between coastal and midland regions with intervening promontories of the crystalline rocks. The Quilon Formation is mainly seen at Paravur ( $08^{\circ}48'00''$ :  $76^{\circ}40'00''$ ) Padappakkara ( $08^{\circ}58'30''$ :  $76^{\circ}38'00''$ ) and some other places around Kollam and Alappuzha districts.

### **Quilon Formation**

The Quilon Formation consisting of fossiliferous shell limestone alternating with thick beds of sandy clays and calcareous clays have been reported from Padappakkara (type locality), Nedumgolam, Edavai ( $8^{\circ}45'20''$ :  $76^{\circ}42'00''$ ) and Varkala ( $8^{\circ}44'00''$ :  $76^{\circ}43'00''$ ) and Cherthala ( $9^{\circ}41'00''$ :  $76^{\circ}20'00''$ ) along the west coast of Kerala. The Quilon limestone contains numerous fossils of foraminifera, corals, echinoids and molluscs. The Lower Miocene age for lower stratigraphic horizons and the Upper Miocene age for the topmost beds of the Quilon Formation indicate the lower and upper age limits of these marine sediments. The

predominance of black clays, sandstone, bluish grey brackish water shell limestone and nodular limestone clearly indicate deposition in a lagoonal condition .

### **Warkalli Formation**

The Warkalli Formation of Mio-Pliocene age extends all along the Kerala coast. The type section of the Warkalli Formation described by King (1882) is from the sea cliff at Varkala. The exposed section at Varkala cliff is 28-30 m thick consisting of unconsolidated sands of variegated clays, white plastic clays, and carbonaceous sandy clays enclosing impersistent seams and lenses of lignite. The carbonaceous clays and lignite are often impregnated with nodules of marcasite.

Fairly thick beds of carbonaceous clays with lignite seams occur around Nadayara kayal, Tamarakulam (9°08': 76°37'), Puliur (9°18'00": 76°35'00"), Payangadi (12°00'20": 75°15'40"), Nileswaram (12°15'00": 75°07'00"), Kanhangad (12°17'40': 75°05'00") and in the cliff sections near Cheruvathur (12°13'00": 75°09'50"). The most characteristic feature of the Warkalli Formation is the impersistent nature of the constituent beds, suggestive of shallow basin margin deposits.

### **Laterite**

Kerala is the home of the laterite as it was first named by the Dutch traveller, Buchanan 1807. Laterite is widespread in its distribution in the midland region of Malappuram, Kannur and Kasaragod districts where it forms well-defined mesas. The Archaean crystalline rocks and the Tertiary sedimentary rocks are extensively lateritised. The laterite has wide areal distribution in the State and occurs at all levels upto 2000 m, height though mostly restricted to an altitude of 50-150 m above MSL. in the coastal and midland region. A few bauxitic patches also occur within the laterites. The thickness of laterite cappings varies from a few metres to 50 metre at places. At Chovvara (8°21'30"; 77°01'30") in Thiruvananthapuram District and Chattannur (8°50'30"; 76°46'30") and Kundara (8°57'00": 76°40'30") in Kollam District, a zone of about 2 m thick bauxite is recognised at the contact between the crystallines and the overlying sedimentary rocks. The overlying sedimentary column is also blanketed by laterite of varying thickness. The bauxite at the base of the sedimentaries indicates an earlier pre-Warkalli spell of lateritisation. Further, the erosional features on the top part of the bauxite horizon corroborates the antiquity of the earlier spell of lateritisation (Mallikarjuna and Kapali, 1980).

Generally, the laterite after the crystalline rocks is compact and the top crust moderately indurated. The dark brown crust passes downward to pink and buff coloured soft laterite. Quartz veins, joints and fractures can be traced from the top to the bottom of the laterite profile. The laterite profile over pyroxene granulites, meta-ultramafites and gneisses are characterised by relict foliation that conforms to those of the subjacent rocks which indicate the *insitu* nature of the laterite. Porous and spongy texture is discernible in laterites, after meta-ultramafites. Laterite after the Tertiary sedimentaries is well indurated at the top for about 2 to 5 m. Downwards, the profile grades into soft laterite with remnants of gritstone and culminates into a zone of variegated clay.

### **Quaternary sediments**

Recent to sub-Recent sediments of coastal sands, sticky black clay with carbonized wood, silty alluvium and lagoonal deposits are observed mostly in the low-lying areas from Kollam (11°27'00": 75°40'30") to Ponnani and between Kannur (11°51'30":75°21'45") and Nileswaram (12°15'30":75°08'16"). Alluvium is observed along the major river valleys. At places, along coastal tracts, there are raised sandy beaches composed of fine grained reddish sandy loam known as "terri" sands. Palaeo-beach ridges alternate with marshy lagoonal clay in the coastal area.

The sandy stretches are widest between Alappuzha (9°30': 76°20') and Kottayam (9°35': 76°31'), upto 25 km inland from the shoreline. The Quaternaries of the coastal plain have been classified into (i) the Guruvayur Formation representing the earlier strandline deposits with an elevation of 5-10 m; (ii) the Viyyam Formation of tidal plain deposits; (iii) Periyar Formation being mainly of fluvial deposits and (iv) the Kadappuram Formation representing the beach deposits (Krishnan Nair, 1989).

A pebble bed is traced in Valapattanam and Taliparamba river banks in Kannur district. It is exposed south of Valapattanam (11°55'30": 75 °21'30"), Kambil maloth (11°58':75 °24'), Morazha (11 °58'30": 75°20'30") and Arathiparamba (12°06'00": 75°15'30"). The size of the pebbles ranges in dimension from 4.5 cm x 3 cm to 7 cm x 3 cm with occasional cobbles of size 13 cm x 12 cm. The base of the pebble bed is generally 20 to 40 m above MSL and at places, the pebble bed directly rests over the basement rocks. The pebbles are mostly of quartz and rarely of granite and pyroxene granulite. The distribution of the pebble bed along the major river banks demonstrate it to be flood plain deposits, probably of early Quaternary period (Nair *et al*, 1976). In Malappuram and Kozhikode districts, the pebble bed is traced in the riverine

terraces at Mavur (11°17'45":75°59'00" ), Cheruvannur (11°12'8": 75°49'35") and Chellepparambu (11°14'30":75°59'00"). In Thiruvananthapuram District, the Quaternary pebble bed occurs at an elevation of 45 to 50 m above MSL at Pothenkode (8°37'00": 76°48'56"), Idaikode (8°40'11":76°50'49"), Attingal (8°41'49": 76°48'56") and Andoorkonam (8°36'00": 76°52'30").

Submerged upright tree trunks have been reported from a number of places in the coastal area of Kottayam and Alappuzha districts, indicating neotectonic reactivation in the area. Carbon dating of a sample from the submerged forest at Iravimangalam indicate an age of 7050 ± 130 B.P (Pawar *et al*, 1983).

### **Structure**

The structural grain of the southern Peninsula is controlled mainly by the NNW-SSE trending near longitudinal Dharwarian trend which had folded all earlier structures. Since Kerala State falls in the western limb of the mega-structure almost all the rock distribution is aligned in NW-SE direction. However, detailed structural studies carried out in selected parts of the Kerala (Nair and Nair, 2001) had shown that (a) the earliest folds (F<sub>1</sub>) which are represented both on mesoscopic and megascopic scale are tight appressed folds of asymmetrical nature which had given rise to axial plane foliations with characteristic platy mineral alignments (b) the F<sub>2</sub> folds on these foliations (post-folial) are open symmetrical and have developed mainly on megascopic scale and control the disposition of the major lithologies. (c) Subsequent folds (F<sub>3</sub>) which deform F<sub>1</sub> and F<sub>2</sub> axial plane traces are broad folds on mega-scale identified with the longitudinal Dharwarian trends and (d) a broad swerve on these Dharwarian trends in ENE-WSW is also decipherable (Fig.2).

Detailed analysis of the remote sensing data had revealed the presence of a number of significant lineament patterns in WNW-ESE, NW-SE, NNW-SSE, NNE-SSW and ENE-WSW directions (Nair, 1990). Mega and intermediate lineaments in WNW-ESE were originally crustal fractures and shears which got sealed or obliterated by a number of igneous emplacements of alkali granite, syenite, gabbro, anorthosite, granophyre etc. The emplacements along the Bavali lineament and those along the Achenkovil lineament both of which trending in this direction had given ages ranging from 500 – 678 Ma. Hence they are identified to be the oldest lineament. The Bavali lineament forms the western termination of the Moyar shear. The NW-SE trending lineaments constitute mega lineaments and coincide with the basic dykes occurring throughout the length and breadth of the state. These dykes have

given ages ranging from 61 to 144 Ma. The NNW-SSE trending lineaments are generally intermediate lineaments and are attributed to fractures, faults and major joint patterns in the area. It is recognized that the NNW-SSE trending lineaments define a weak zone along which the west coast evolved by faulting. The eastern limit of the Tertiary basin is found restricted along this lineament direction. These lineaments occurring along the west coast are be active as suggested by the progradation of the coast west of these lineaments (Nair, 1987). The lineaments in NNE-SSW are prominent and are identified with major fractures and this together with those in NNW-SSE are taken to constitute a conjugate system of faults in a N-S compressive regime due to the collision of the Indian plate. The ENE-WSW trending lineaments are intermediate lineaments and are well- developed in the northern parts of the Kerala . Since these lineaments truncate other lineaments as evidenced especially in the coastal stretches it is considered the youngest. Many a recent tremors reported are aligned in this direction and hence considered neotectonically active.

### **Metamorphism**

The Precambrian crystalline rocks of Kerala are chiefly metapelites, charnockites with associated gneisses and granulites, schistose rocks with distinct metapelitic and metamafic / ultramafic affinity and granitic derivatives which include the Peninsular gneisses and migmatites. Except the Wynad schists and the Vengad group, the bulk of the crystalline rocks show granulite to upper amphibolite facies of metamorphism. Wynad schist displays a prograde amphibolite facies metamorphism and the retrogression of these rocks leads to lower amphibolite facies metamorphism. The vast charnockite belt occurring on either side of the Wynad schist belt, in north Kerala, shows petrographic evidences of prograde and retrograde reactions (Nambiar, 1996). The rocks of the Vengad Group show greenschist to lower amphibolite facies of prograde metamorphism. The older intrusive bodies show effects of incipient metamorphism, marked by clouding of feldspar and bending of twin lamellae.

Recent investigations on the pressure – temperature range for the formation of characteristic mineral suits within the metamorphic rocks provide a fair idea on the poly-metamorphic history of the rock suits. Rocks of the Khondalite belt of south Kerala indicate a temperature range of 650 to 850°C and pressures 5 to 6 kb (Srikantappa *et al*, 1985). In the Thiruvananthapuram area, the temperature at the peak of metamorphism indicated by the mineral assemblages of the calc-silicate rocks is about 830°C at 5 K bar considering the vapour absent garnet forming equilibria (Satish Kumar and Santosh, 1996). The scapolite equilibria indicates a peak metamorphic temperature of above 800°C. Stable isotopes in the marble bands suggest that there was no

pervasive infiltration of external fluids. Local infiltration of external carbonic fluid took place during decomposition. Synthesis of such data from different lineament/shear bound segments in Kerala indicate varying metamorphic conditions and uplift history. It is also summarised that there is a progressive decline in the uplift of different segments from north to south (Soman, 1997).

### **Economic minerals**

The existing level of information on the mineral resources of the State indicate rich deposits of heavy mineral sands along the coast and rich clay deposits. Other significant occurrences include bauxite, iron ore, graphite, limestone and gold.

### **Major minerals**

They include china clay, Bauxite/laterite, Ilmenite, Rutile, Zircon, Sillimanite, Leucoxene, Monazite, Silica Sand, Lime Shell, Limestone, Quartz, Graphite

### **Minor minerals**

Granite Building stone, Granite dimension stone, Laterite, River sand, Ordinary sand, lime shell, ordinary earth, brick clay



 **SUSTAINABLE DEVELOPMENT GOALS**

